

BMB Reports – Manuscript Submission

Manuscript Draft

Manuscript Number: BMB-17-049

Title: Fine-tuning of gene expression dynamics by the Set2-Rpd3S pathway

Article Type: Perspective (Invited Only)

Keywords: Set2; H3K36 methylation; Rpd3S HDAC; histone deacetylation; Gene expression dynamics

Corresponding Author: TaeSoo Kim

Authors: Bo Bae Lee¹, Ji Hyun Kim¹, TaeSoo Kim^{1,*}

Institution: ¹Department of Life Science and the Research Center for Cellular Homeostasis, Ewha Womans University, Seoul 03760, Korea,

Manuscript Type: Perspective

Fine-tuning of gene expression dynamics by the Set2-Rpd3S pathway

Bo Bae Lee¹, Ji Hyun Kim¹, TaeSoo Kim*

Department of Life Science and the Research Center for Cellular Homeostasis,
Ewha Womans University, Seoul 03760, Korea.

Corresponding Author's Information: tskim@ewha.ac.kr

¹These authors equally contributed this work.

Keywords: Set2, H3K36 methylation, Rpd3S HDAC, histone deacetylation, Gene expression dynamics

Running title: The Set2-Rpd3S HDAC optimizes gene expression dynamics

Kim et al., (2016) Nature Communications 7:13534, DOI: 10.1038/ncomms13534

ABSTRACT

RNA polymerase II-interacting the Set2 methyltransferase co-transcriptionally methylates histone H3 at lysine 36 within the body of genes. This modification facilitates histone deacetylation by Rpd3S HDAC in 3' transcribed regions to suppress cryptic initiation and slow elongation. Although this pathway is important for global deacetylation, no strong effects on genome-wide transcription have been seen under optimized laboratory conditions. In contrast, this pathway slows the kinetics of mRNA induction when target genes are induced upon environmental changes. Interestingly, a majority of Set2-repressed genes are overlapped by lncRNA transcription that targets H3K36 methylation and deacetylation by Rpd3S HDAC to mRNA promoters. Furthermore, this pathway delays the induction of many cryptic transcripts upon environmental changes. Therefore, the Set2-Rpd3S HDAC pathway functions to fine-tune expression dynamics of mRNAs and ncRNAs.

Chromatin, a complex of DNA and histones plays important roles in many processes including RNA polymerase II (RNA polII) transcription, DNA replication, and DNA repair. Nucleosome, the basic repeating unit of chromatin consists of four core histones, H3, H4, H2A, and H2B and 147 base pairs of DNA. Post-translational modifications of histones including acetylation, methylation, phosphorylation, and ubiquitination positively or negatively affect RNA polII transcription by regulating local nucleosome structure. Histone acetylation is known to disrupt the interaction between DNA and histone proteins and highly correlated with transcription frequency. This modification is dynamically regulated by both histone acetyltransferases (HATs) and histone deacetylases (HDACs). Histone methylation also affects local histone acetylation by acting as a binding site for either HATs or HDACs.

The Set2 methyltransferase preferentially interacts with phosphorylated CTD at both serine 5 and serine 2 of Rpb1, the largest subunit of RNA polII and methylates histone H3 at lysine 36 within the body of genes. H3K36 methylation by Set2 is involved in RNA polII elongation and co-transcriptional RNA processing. Since nucleosomes within the body of genes directly block elongation, histones need to be acetylated by HATs and evicted to promote elongation. However, once RNA polII transcribes through the gene, histones must be deacetylated to re-establish a repressive nucleosome structure and to suppress spurious transcription within the genes. Set2-targeted H3K36 methylation facilitates histone deacetylation by Rpd3S (Rpd3 small) HDAC within coding regions. Two subunits, Eaf3 and Rco1, are important for binding of this complex to chromatin. Eaf3 chromodomain preferentially binds to methylated histones on H3K36 and Rco1 PHD finger interacts with unmodified histone tails. Both interactions stimulate histone deacetylation by Rpd3S HDAC. Although previous studies on the Set2-Rpd3S HDAC pathway revealed molecular mechanisms for how the pathway is targeted and inhibits transcription from cryptic promoters, its function in mRNA transcription remains elusive. Interestingly, global gene expression

pattern was not strongly affected in mutants for this pathway under optimized growth conditions.

To further understand the function of the Set2-Rpd3S HDAC pathway in mRNA regulation, we analyzed genome-wide transcript levels in wild type and *SET2* deleting cells during carbon source shifts. Although steady-state levels of transcripts were not changed, ~80 genes showed a more rapid induction in the absence of Set2. Furthermore, we also identified 757 cryptic promoters repressed by Set2. Interestingly, 416 cryptic promoters not reported previously were induced by distinct carbon sources. These findings suggest that the Set2-Rpd3S HDAC pathway plays important roles in maintaining optimal expression dynamics of mRNAs and non-coding RNAs(ncRNAs) upon environmental changes (**Figure 1A**).

Interestingly, a majority of Set2-repressed genes had an overlapping long non-coding RNA(lncRNA) transcription from an upstream or an antisense promoter. Furthermore, these genes showed non-canonical patterns of histone modifications. Although H3K4me3 and H3K36me3 are in general enriched at promoters and 3' transcribed regions, respectively, Set2-repressed mRNA promoters showed high levels of H3K36me3 but not H3K4me3. We found that H3K36me3 at mRNA promoters targeted by an overlapping lncRNA transcription induced histone deacetylation by Rpd3S HDAC to delay the induction of target mRNA genes (**Figure 1A**). A recent study from our group has revealed that H3K4me2-Set3 HDAC pathway also requires an overlapping lncRNA transcription to modulate the induction of target genes (Kim et al., (2012) *Cell*, 150, 1158-69. DOI: 10.1016/j.cell.2012.08.016). Although the genes regulated by both H3K4me2-Set3 HDAC and Set2-Rpd3S HDAC pathways tend to be overlapped by a lncRNA transcription, the two pathways affect the activation of mostly distinct target genes. An interesting difference between Set3- or Set2-

repressed genes is the distance between mRNA and lncRNA promoters (**Figure 1B**). Surprisingly, alteration of the distance between the mRNA and lncRNA promoters switches gene repression by the Set2-Rpd3S to H3K4me2-Set3 HDAC. Our findings suggest that although lncRNA can be transcribed from any site of a genome, locations of overlapping lncRNA promoters may be important for targeting distinct histone methylations and chromatin regulators to promoter regions of an associated mRNA gene (**Figure 1B**).

Whereas gene-specific transcriptional activators or repressors primarily turn on or off their target genes, chromatin regulators appear to function as modulators that fine-tune the kinetics of gene induction or repression. This type of gene regulation seems to be important for cells growing their natural conditions as they must reprogram gene expression patterns to adapt to rapidly changing environmental conditions. Therefore, fine-tuning of gene expression dynamics by chromatin regulators is likely important for normal development processes, cellular adaptation, and proper responses to environmental stresses.

Acknowledgements

This work was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIP) (No. 2012R1A5A1048236).




Figure 1. Models for regulation of promoter dynamics by the Set2-Rpd3S pathway.

A. IncRNA transcription from an upstream targets H3K36me3 and Rpd3S HDAC to mRNA promoters. Histone deacetylation at mRNA promoters by Rpd3S HDAC delays the kinetics of gene induction upon environmental changes. Transcription from the mRNA promoter of a gene targets H3K36me3 and Rpd3S HDAC to inducible cryptic promoters. The resulting deacetylation by Rpd3S HDAC slows the rate of cryptic RNA induction.

B. Location of the IncRNA promoter determines the repressive effect of Set3 HDAC and Rpd3S HDAC. H3K4me2 and H3K36me3 peak at 5' and 3' regions of IncRNA transcription, respectively. Set3 HDAC deacetylates H3K4me2-enriched nucleosomes to repress the

mRNA promoter in 5' regions. In 3' regions of lncRNA transcription, Rpd3S HDAC deacetylates histones and represses mRNA promoters.