

BMB Reports – Manuscript Submission

Manuscript Draft

**Manuscript Number:** BMB-19-097

**Title:** Neuronal function and dysfunction of CYFIP2: from actin dynamics to early infantile epileptic encephalopathy

**Article Type:** Mini Review

**Keywords:** cytoplasmic FMR1-interacting protein 2; WAVE regulatory complex; actin polymerization; neuronal synapse; early infantile epileptic encephalopathy

**Corresponding Author:** Kihoon Han

**Authors:** Kihoon Han<sup>1,\*</sup>, Yinhua Zhang<sup>1</sup>, Yeunkum Lee<sup>1</sup>

**Institution:** <sup>1</sup>Department of Neuroscience and <sup>2</sup>Department of Biomedical Sciences, College of Medicine, Korea University, Seoul 02841, South Korea,

**Manuscript Type:** Mini Review

**Title:**

Neuronal function and dysfunction of CYFIP2:  
from actin dynamics to early infantile epileptic encephalopathy

**Author's name:** Yinhua Zhang<sup>1,2</sup>, Yeunkum Lee<sup>1,2</sup> and Kihoon Han<sup>1,2,\*</sup>

**Affiliation:**

<sup>1</sup>Department of Neuroscience, College of Medicine, Korea University, Seoul 02841,  
South Korea

<sup>2</sup>Department of Biomedical Sciences, College of Medicine, Korea University, Seoul  
02841, South Korea

**Running Title:** Neuronal function and dysfunction of CYFIP2

**Keywords:** cytoplasmic FMR1-interacting protein 2 (CYFIP2), WAVE regulatory complex (WRC), actin polymerization, neuronal synapse, early infantile epileptic encephalopathy (EIEE)

**\*Corresponding Author's Information:**

Tel: +82-2-2286-1390; E-mail: neurohan@korea.ac.kr

## ABSTRACT

The cytoplasmic FMR1-interacting protein family (CYFIP1 and CYFIP2) are evolutionarily conserved proteins originally identified as binding partners of the fragile X mental retardation protein (FMRP), a messenger RNA (mRNA)-binding protein whose loss causes the fragile X syndrome. Moreover, CYFIP is a key component of the heteropentameric WAVE regulatory complex (WRC), a critical regulator of neuronal actin dynamics. Therefore, CYFIP may play key roles in regulating both mRNA translation and actin polymerization, which are critically involved in proper neuronal development and function. Nevertheless, compared to CYFIP1, neuronal function and dysfunction of CYFIP2 remain largely unknown, possibly due to the relatively less well established association between *CYFIP2* and brain disorders. Despite high amino acid sequence homology between CYFIP1 and CYFIP2, several *in vitro* and animal model studies have suggested that CYFIP2 has some unique neuronal functions distinct from those of CYFIP1. Furthermore, recent whole-exome sequencing studies identified *de novo* hot spot variants of *CYFIP2* in patients with early infantile epileptic encephalopathy (EIEE), clearly implicating CYFIP2 dysfunction in neurological disorders. In this review, we highlight these recent investigations into the neuronal function and dysfunction of CYFIP2, and also discuss several key questions remaining about this intriguing neuronal protein.

## INTRODUCTION

Actin cytoskeleton dynamics are critically involved in various aspects of neuronal development and function, including early migration, dendritic and axonal outgrowth and branching, and synapse formation and plasticity (1-3). Consistently, several mutations of genes encoding key regulators of the neuronal actin cytoskeleton have been associated with numerous neurodevelopmental and neuropsychiatric disorders, such as autism spectrum disorders (ASDs), intellectual disability (ID), schizophrenia (SCZ), and bipolar disorder (4-6). Among the key regulators, the WAVE regulatory complex (WRC) is a ~400 kDa heteropentameric protein complex consisting of the Wiskott–Aldrich syndrome protein family verprolin-homologous protein (WAVE), cytoplasmic FMR1-interacting protein (CYFIP), Nck-associated protein (NAP), Abelson-interacting protein (ABI), and hematopoietic stem progenitor cell 300 (HSPC300) (7, 8). The WRC is basally inactive toward its downstream actin-nucleator, i.e., the actin-related protein 2/3 (Arp2/3) complex, because the activity-bearing C-terminal VCA (verprolin homology, central, and acidic regions) domain of WAVE is sequestered by an intermolecular interaction with CYFIP (8). However, diverse cellular signaling can release the VCA domain from the CYFIP, enabling it to bind to and activate the Arp2/3 complex. Specifically, inositol phospholipids and the active form of the small GTPase Rac1 (Rac1-GTP) can cooperatively recruit the WRC to the plasma membrane and induce its conformational changes (8). Considering that Rac1-GTP directly binds to the CYFIP during the activation process, CYFIP is involved in both basal inhibition and Rac1-induced activation of the WRC. Furthermore, phosphorylation of several residues of WAVE, CYFIP, and ABI can also modulate the WRC activity (9-11).

Two evolutionarily highly conserved members of the *CYFIP* gene family, namely

*CYFIP1* and *CYFIP2* (also referred to as *SRA-1* and *PIR121*, respectively) exist (12). They are ~145 kDa proteins, with a high homology in their amino acid sequences (88% identity and 95% similarity), suggesting similar functions at the molecular level. Originally, the *CYFIP1/2* proteins were identified as binding partners of the fragile X mental retardation protein (FMRP) (12), whose loss causes the fragile X syndrome (FXS), which is characterized by ID, autistic behaviors, and dysmorphic features (13). Indeed, in addition to the WRC, *CYFIP1* forms a complex with both the FMRP and the eukaryotic initiation factor 4E (eIF4E) to regulate the translational initiation of target messenger RNAs (mRNAs) (14). Notably, the brain-derived neurotrophic factor (BDNF)-induced activation of Rac1 has been shown to redistribute *CYFIP1* from the FMRP-eIF4E complex to the WRC, thereby orchestrating both activity-dependent local protein synthesis and actin remodeling in neuronal synapses (15). Clinically, structural abnormalities (deletions and duplications) of the chromosomal region (15q11-q13) encompassing *CYFIP1* were identified in patients with ASDs and ID (16). *CYFIP1* has also been associated with SCZ (17). Furthermore, several lines of *Cyfp1* mutant mice expressed abnormal synaptic morphology and function, as well as either FXS- or ASD-like behaviors (18-20).

In contrast to *CYFIP1*, the neuronal function and dysfunction of *CYFIP2* remain largely unknown, possibly given the relatively less well established association between *CYFIP2* and brain disorders thus far (16). Nevertheless, despite the high sequence homology between *CYFIP1* and *CYFIP2*, several lines of evidence suggest that they may have distinct functions *in vivo*. For example, both *Cyfp1* and *Cyfp2* null mice display lethality at different developmental time points (i.e., at early embryonic and perinatal stages, respectively) (20-23). Furthermore, recent whole-exome sequencing (WES)

studies identified *de novo* hot spot mutations of *CYFIP2* in patients with early infantile epileptic encephalopathy (EIEE) (24, 25). In this review, we highlight recent *in vitro*, animal model, and human genetic investigations revealing the neuronal function and dysfunction of *CYFIP2*, and discuss the key remaining issues for a better understanding of this intriguing, yet largely uncharacterized, protein in the nervous system.

### ***IN VITRO* NEURONAL STUDIES**

Through the overexpression of green fluorescent protein (GFP)-tagged *CYFIP1* and *CYFIP2*, Pathania et al. investigated the subcellular localization of these proteins in cultured rat hippocampal neurons (26). Both *CYFIP1*-GFP and *CYFIP2*-GFP exhibited punctate clusters highly localized to the dendritic spines, which are small protrusions on the neuronal dendrites representing excitatory postsynaptic compartments in mature neurons (27). Consistently, both proteins co-localized with endogenous excitatory synaptic markers, namely the Homer protein homolog (Homer) and the vesicular glutamate transporter (VGLUT). Indeed, the excitatory synaptic localization of the *CYFIP1/2* proteins was further supported by several biochemical studies. For example, the *CYFIP1/2* proteins were identified as components of the postsynaptic proteome through mass-spectrometry analyses of both the purified human and rodent postsynaptic density (28, 29). Moreover, both proteins were recognized as key nodes of the mouse *in vivo* postsynaptic protein interactome (30), whereas *CYFIP2* was identified as an *in vivo* postsynaptic scaffold SH3 and multiple ankyrin repeat domains 3 (Shank3) interactor (31). In addition to their postsynaptic localization, Pathania et al. also reported that overexpression of either *CYFIP1* or *CYFIP2* increased dendritic branching and outgrowth of cultured hippocampal neurons (26). In the same study, the effects of *Cyfip1*, but not of

*Cyfp2*, haploinsufficiency (in *Cyfp1*<sup>+/-</sup> mice) on both hippocampal dendritic spine density and morphology were explored, and an increase in immature long and thin spines compared to wild-type (WT) mice was observed.

A large-scale *in vivo* phospho-proteomic analysis of nine mouse tissues (32) identified two brain-specific phosphorylation sites (S582 and T1067) of CYFIP2 (11). Furthermore, Lee et al. overexpressed WT, phospho-blocking (i.e., S582A and T1067A), and phospho-mimetic (i.e., S582D and T1067E) mutants of CYFIP2 in cultured hippocampal neurons to understand functional significance of such sites (11). With regard to dendritic spine regulation, overexpression of the T1067A mutant, but not that of either WT or other mutants, decreased the presence of stubby spines compared to control neurons, suggesting a dominant-negative effect of T1067A. In addition, while overexpression of WT CYFIP2 increased neurite outgrowth, the same was not true for T1067A. Intriguingly, when mapped on the crystal structure of the WRC (31), CYFIP2 T1067 was positioned at the interaction interface between the CYFIP and the NAP, and T1067 phosphorylation was predicted to weaken such an interaction (11). Lee et al. proposed a model based on these findings. It described that CYFIP2 T1067 phosphorylation, together with additional modifications of the WRC (such as bindings of inositol phospholipids and Rac1-GTP (8)), contributed to WRC activation through the disassembling of the self-inhibitory interactions among its components (11). In contrast to T1067 phosphorylation, the functional roles of CYFIP2 S582 phosphorylation, which is known to occur on the N-terminal outer surface of CYFIP2 in the WRC, are yet to be defined (11).

## ANIMAL MODEL STUDIES

The *in vivo* neuronal functions of CYFIP were initially characterized in the *Drosophila*, which expresses only one homolog of CYFIP (*dCYFIP*) (16). In fact, Schenck et al. demonstrated that CYFIP is specifically expressed in the central nervous system (CNS) of *Drosophila*, and is localized at synaptic terminals in the neuromuscular junctions (NMJs) (33). The length of the synaptic terminal was reduced in the NMJs of *dCYFIP* mutant larvae, compared to WT larvae, while the synaptic boutons had immature supernumerary buds (33). Notably, the reduced synaptic terminal length of *dCYFIP* mutants was the opposite phenotype of that observed in the *Drosophila FMR1* (*dFMR1*) null NMJs (34). Indeed, Abekhoukh et al. determined that the NMJs of *dCYFIP;dFMR1* double homozygous mutants were similar to WT NMJs, and not to those present in either of the single mutants. This suggests a genetic interaction between *dCYFIP* and *dFMR1* associated with the regulation of NMJ synaptic structures (35). Furthermore, Zhao et al. identified further detailed structural and functional changes in *dCYFIP* mutant NMJs (36). Specifically, enlarged synaptic vesicles and additional cisternae in the synaptic boutons were found through electron microscopic analysis of *dCYFIP* mutants. Moreover, given that *dCYFIP* mutants also showed functional defects related to the release of neurotransmitters under high-frequency stimulation, Zhao et al. suggested that dCYFIP regulates synaptic endocytosis by inhibiting F-actin assembly (36).

The *nevermind* (*nev*) mutant zebrafish was isolated from a large-scale forward genetic screening to identify the genes involved in retinotectal axon pathfinding (37, 38). Specifically, the dorsonasal axons of retinal ganglion cells (RGCs) are missorted with ventral axons in *nev* mutant zebrafish, and are abnormally projected through the dorsal tectum (even though they still targeted to the ventral tectum). Pittman et al. identified the *nev* gene to encode zebrafish *Cyfip2* (39). In fact, while the zebrafish expresses both



CYFIP1 and CYFIP2 (86% identical), its CYFIP2 is 98% identical to that in humans (12). Pittman et al. also reported a broad *Cyfp2* expression in the zebrafish CNS, even though the other axonal tracts of *nev* mutants, including those of both the primary motor neurons and Mauthner neurons, were normal (39). Furthermore, Pittman et al. found that retinal lamination was disrupted in *nev* mutants (39). Additionally, Cioni and colleagues identified more detailed molecular and cellular mechanisms behind the optic tract missorting in *Cyfp2* mutant zebrafish (40). By combining the molecular replacement approach and time-lapse imaging of axonal growth cones, they showed the involvement of CYFIP2 in both homotypic axonal fasciculation and heterotypic axonal repulsion. Specifically, CYFIP2 was translocated to the growth cone in response to axon-axon contact, and regulated the filopodial dynamics to induce either axonal fasciculation or repulsion. Unlike *Cyfp2*, CRISPR/Cas9-mediated deletion of *Cyfp1* caused axonal growth defects in the optic tract, suggesting that CYFIP1 and CYFIP2 are specifically involved in axonal growth and sorting, respectively (40). In addition to the visual system, the CYFIP2 function in the auditory system of the zebrafish was also characterized. Marsden et al. performed an n-ethyl-n-nitrosourea (ENU)-based forward genetic screening to identify zebrafish mutants with reduced auditory startle thresholds (i.e., enhanced startle response) and isolated the *Cyfp2* mutant as one of five lines having such a phenotype (41). The zebrafish startle circuit is composed of auditory afferents (VIII), excitatory spiral fiber (SF) interneurons, and Mauthner cells. Marsden et al. showed that the activity of this circuit was enhanced in *Cyfp2* mutant zebrafish, as a result of either the increased excitability of SF interneurons or the enhanced excitatory synaptic input to SF interneurons (41). Although the detailed molecular mechanism underlying such a phenomenon is yet to be defined, the FMRP was not shown to be involved in this auditory

phenotype of *Cyfip2* mutant, as the *fmr1* mutant zebrafish presented a normal innate startle threshold.

The initial genetic evidence supporting the neuronal functions of CYFIP2 in mice derived from quantitative trait locus (QTL) analyses aimed at identification of the genetic loci contributing to the behavioral differences between the C57BL/6J and C57BL/6N mouse substrains (21, 42). Specifically, the C57BL/6N substrain was branched out from original C57BL/6J substrain in the 1950s, and is currently the second major C57BL/6 mouse substrain. However, whole genome sequencing identified only approximately 10,000 single nucleotide polymorphisms (SNPs) between the two substrains (43). Therefore, QTL analysis can be useful for detecting the causal genetic factors underlying certain phenotypic differences between such substrains. Kumar et al. described the C57BL/6N substrain as having a lower behavioral response (as measured by locomotor activity) to psychostimulants (such as cocaine and methamphetamine), compared to C57BL/6J (21). Through the performance of a QTL analysis, Kumar et al. identified that a nonsynonymous SNP (G to A at the 46,036,117 base pair of chromosome 11, mm9) producing a serine-to-phenylalanine missense mutation (S968F) in CYFIP2 was responsible for the difference in the psychostimulant responses (21). Successively, Kirkpatrick et al. used a similar QTL analysis to show that such a *Cyfip2* variant was also causally associated with the enhanced binge eating-like behavior of the C57BL/6N substrain, compared to the C57BL/6J substrain (42). With regard to the molecular mechanism behind such a phenomenon, Kumar et al. reported the S968F CYFIP2 proteins to be less stable than WT CYFIP2 when overexpressed in HEK293T cells (21). However, recent structural analysis of a WRC-Rac1 complex by cryo-electron microscopy suggested that the S968F CYFIP2 could cause excessive activation of the WRC by Rac1

(44). Therefore, further molecular and cellular studies are required to understand the process by which a single CYFIP2 variant changes the behavioral phenotypes in mice.

Both *Cytip1* and *Cytip2* null mice show lethality at different developmental time points. Specifically, while the *Cytip1* null embryos die around embryonic day 9.5 (E9.5) (20, 26), the *Cytip2* null (*Cytip2*<sup>-/-</sup>) mice die perinatally (21, 22). Recently, Zhang et al. revealed that, although *Cytip2*<sup>-/-</sup> mice survived until E18.5, with an expected Mendelian inheritance ratio they would all die at postnatal day 0 (P0) (23). Therefore, it is conceivable that *Cytip2*<sup>-/-</sup> mice could die very soon after birth (21). Although the detailed cause of lethality remains unknown, Zhang et al. also reported that *Cytip2*<sup>-/-</sup> mice were smaller than either WT or heterozygous (*Cytip2*<sup>+/-</sup>) littermates at E18.5 (23). Specifically, the crown-rump length was shorter in *Cytip2*<sup>-/-</sup> mice, suggesting defects in their body curvature. Nevertheless, both the brain size and overall cytoarchitecture were comparable between WT, *Cytip2*<sup>+/-</sup> and *Cytip2*<sup>-/-</sup> mice at E18.5. Subsequent to an RNA-sequencing analysis, however, Zhang et al. found the extracellular matrix (ECM)-related genes to be significantly altered in the *Cytip2*<sup>-/-</sup> embryonic cortex, compared with the WT cortex (23). Whether the abnormal expression of ECM-related genes can affect the neuronal development and function in *Cytip2*<sup>-/-</sup> embryos, and whether such changes can be associated with their early postnatal lethality remains to be investigated.

Given the early postnatal lethality of *Cytip2*<sup>-/-</sup> mice, *Cytip2*<sup>+/-</sup> mice were used to characterize the functions of CYFIP2 in adult brains. Han et al. hypothesized that CYFIP2 could be one of the key actin-regulatory proteins mediating the dendritic spine abnormalities observed in fragile X mental retardation 1 (*Fmr1*)-null (*Fmr1*<sup>-/-</sup>) mice, an animal model of FXS (22). In addition to the protein-protein interaction between CYFIP2 and FMRP (12), the *Cytip2* mRNA has been ranked as the ninth FMRP target among 842

FMRP-associated brain polyribosomal mRNAs (45). Therefore, it is plausible that the basal and/or activity-induced translation of *Cyfp2* mRNA can be altered in *Fmr1*<sup>-/-</sup> mice, possibly contributing to the abnormal actin-rich dendritic spines found in these mice. Moreover, Han and colleagues found that, similar to *Fmr1*<sup>-/-</sup> mice, adult *Cyfp2*<sup>+/-</sup> mice exhibited some FXS-like behaviors, including hyper locomotor activity, reduced acoustic startle response, and enhanced pre-pulse inhibition (22). Notably, such behavioral phenotypes were aggravated in the *Fmr1*<sup>-/-</sup>;*Cyfp2*<sup>+/-</sup> double mutant mice. In concordance, the dendritic spine abnormality seen in the *Fmr1*<sup>-/-</sup> cortex (i.e., the increase in immature thin spines) was aggravated in double mutant mice (22). The basal levels of both the CYFIP2 and WAVE1 proteins were normal in the hippocampus and cortex of *Fmr1*<sup>-/-</sup> mice. Han et al. showed the treatment of cultured cortical neurons with a metabotropic glutamate receptor (mGluR) agonist 3,5-dihydroxyphenylglycine (DHPG) to induce the expression of CYFIP2 proteins, which was blocked by the translational inhibitor cycloheximide (22). Notably, both the DHPG-induced expression of CYFIP2 and the DHPG-induced dendritic spine remodeling were impaired in both *Fmr1*<sup>-/-</sup> and *Cyfp2*<sup>+/-</sup> cortical neurons. Therefore, *Cyfp2* mRNA can be translated in an mGluR-dependent manner, as a target of the FMRP-associated polyribosome, and newly expressed CYFIP2 proteins may be involved in dendritic spine remodeling (22). Nevertheless, the causal relationship between such molecular and cellular changes in *Cyfp2*<sup>+/-</sup> neurons and the behavioral phenotypes of *Cyfp2*<sup>+/-</sup> mice remains to be further validated.

## **CYFIP2 MUTATIONS IN EARLY INFANTILE EPILEPTIC ENCEPHALOPATHY**

In contrast to the well-established associations between *CYFIP1* and either ASDs, ID, or

SCZ, *CYFIP2* has not been clearly associated with any brain disorders until recently (16). Although interstitial deletions of the chromosomal region 5q33.3-q35.1 harboring *CYFIP2* have been identified in individuals with developmental delay, mild to severe ID, and seizures (46, 47), those deletions included tens of additional genes, preventing the identification of the causal gene(s). Furthermore, reduced protein expression of *CYFIP2* in postmortem brains of patients with SCZ (48) and Alzheimer's disease (49) was reported, even though its causal relationship with the disease pathophysiology was not clearly established. Notably, two recent WES studies similarly identified *de novo* hot spot (Arg87) variants in *CYFIP2* from patients with EIEE (24, 25). EIEE (OMIM 308350, also referred to as the Ohtahara syndrome) is a neurological disorder characterized by the occurrence of seizures, which begin within a year of birth (50). This disorder often progresses to West syndrome, which is characterized by epileptic spasms with hypsarrhythmia on the electroencephalogram (EEG) and severe psychomotor developmental delay (51).

In the first study, Nakashima et al. identified three *de novo* *CYFIP2* variants at the Arg87 residue (Arg87Leu, Arg87Cys, and Arg87Pro) in four unrelated patients diagnosed with West syndrome (24). All four individuals presented their first seizure event prior to the age of 6 months and reported additional common features, including hypotonia, microcephaly, developmental delay, and ID. Using the available crystal structure of the WRC (8), Nakashima and colleagues described the Arg87 residue of *CYFIP1* (the solved WRC structure contains *CYFIP1*, not *CYFIP2*) to be at an interface between *CYFIP1* and *WAVE1* (24). Specifically, the side chain of Arg87 formed hydrogen bonds with both the main chain of Tyr151 in *WAVE1* and the side chains of Glu625 and Glu690 in *CYFIP1*. Importantly, the three Arg87 variants were predicted to disrupt these hydrogen bonds, thereby destabilizing the interaction between *CYFIP1* and *WAVE1*. Indeed, all three

CYFIP2 variants showed weaker interaction with the purified VCA domain of the WAVE, compared to WT CYFIP2 (24). Furthermore, transfection of CYFIP2 mutants, but not WT, induced an aberrant F-actin accumulation in cultured mouse melanoma cells (24); an expected finding, given that the mutants released the VCA domain of the WAVE, leading to activation of the downstream Arp2/3 complex. In a subsequent investigation, Peng et al. performed a WES analysis of 56 Chinese families with West syndrome and identified 112 variants in 89 genes, which included a *de novo* Arg87Cys variant of *CYFIP2* (25). Intriguingly, an in-frame deletion variant of *CYFIP1* was also described. Overall, these studies suggest that *de novo* hot spot (Arg87) variants of *CYFIP2* can be causally associated with West syndrome, potentially through the induction of aberrant WRC activation (i.e., gain-of-function) and F-actin polymerization.

## PERSPECTIVES AND CONCLUSIONS

As highlighted in the current review, although a growing amount of evidence supports distinct *in vivo* functions of CYFIP1 and CYFIP2 (35, 40), the detailed mechanisms underlying such a phenomenon are largely unknown. Distinct spatiotemporal expression patterns of CYFIP1 and CYFIP2 may contribute to this (40, 52). For example, while a preliminary analysis found both the *Cyfp2* mRNA and protein to be enriched in the cortex, compared with other brain regions of mice, those of *Cyfp1* were observed to be evenly expressed in different brain regions (unpublished data). This may, at least partly, explain the cortical, but not hippocampal, dendritic spine changes seen in *Cyfp2*<sup>+/-</sup> mice (22). Moreover, a different expression of CYFIP1 and CYFIP2 in brain non-neuronal cells cannot be excluded. Indeed, a mouse brain cell-type-specific RNA-sequencing database (<http://www.brainrnaseq.org/>) (53) highly detected the presence of *Cyfp1* mRNA in

microglia/macrophage, whereas *Cytip2* mRNA was abundant in newly-formed oligodendrocytes and neurons. In addition to their basal expression, neuronal activity-induced expression of *Cytip1* and *Cytip2* can be differentially regulated. For example, while *Cytip2* mRNA was identified as a high ranked FMRP target (45), the same was not true for *Cytip1*, indicating that only the dendritic transport, stability, and local translation of *Cytip2* mRNA can be regulated by FMRP in an activity-dependent manner (22, 54). In accord, *Cytip2* mRNA was detected in neuronal dendrites (55). It is also possible that, despite their high sequence homology, CYFIP1 and CYFIP2 may have distinct groups of protein interactors. For example, in the original study which identified both CYFIP1/2 as direct binding partners of FMRP (12), CYFIP2, but not CYFIP1, was also shown to interact with FMRP-related proteins (FXR1P/FXR2P). Several independent investigations, using different biochemical approaches, recently identified the protein interactors specific to both CYFIP1 and CYFIP2 (15, 21, 30, 40). A side-by-side comprehensive analysis of these protein lists may provide another important insight into understanding the diverse functions of CYFIP1 and CYFIP2 *in vivo*.

Detailed molecular and cellular mechanisms underlying the processes by which the Arg87 variants of *CYFIP2* lead to the neurological symptoms of West syndrome remain to be further investigated (Fig. 1). Nakashima et al. proposed that an aberrant increase in WRC activity and downstream actin polymerization may change the neuronal excitatory synaptic function, thereby disrupting the excitatory/inhibitory (E/I) synaptic balance (24) (Fig. 1A), which is considered one of the key mechanisms behind epilepsy, and other neurodevelopmental and neuropsychiatric disorders (56, 57). Not only postsynaptic functions, but also excitatory (and inhibitory) presynaptic functions may be regulated by CYFIP2 (Fig. 1B), since CYFIP1 regulates presynaptic terminal size and

vesicle release probability in the developing hippocampus (58). Another possibility is that the change in neuronal excitability (neuronal propensity to generate action potential firing) by the *CYFIP2* variants, is caused by affecting the surface expression and/or functional properties of certain ion channels. Notably, Chen et al. indicated that the WRC can potentially interact with ~102 diverse membrane proteins, including protocadherins, G protein-coupled receptors (GPCRs), and ion channels (59). Importantly, a detailed structural analysis revealed that both the CYFIP and ABI of the WRC formed an interaction surface toward those membrane proteins. Therefore, the *CYFIP2* variants found in West syndrome may indirectly affect the interactions between the WRC and certain ion channels, possibly leading in turn to changes in both the ion channel properties and neuronal excitability (Fig. 1C). Furthermore, the FMRP-mediated regulation of expression of the proteins involved in modulating the synaptic E/I balance or the neuronal excitability may also play a role (60). Similarly to CYFIP1, CYFIP2 can form a complex with the FMRP and other RNA-binding proteins to regulate mRNA transport and local translation (40), whereas the Arg87 variants may affect this process (Fig. 1D). An additional comprehensive analysis of the expression, localization, interaction, and activity-dependent regulation of CYFIP2 using various *in vitro* and *in vivo* approaches will elucidate its unique functions, which can have implications for both the proper neuronal development and the pathophysiology of several brain disorders.

## ACKNOWLEDGMENTS

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government Ministry of Science and ICT (NRF-2018R1C1B6001235, NRF-2018R1A6A3A11040508), and by the Brain Research



Program through the NRF funded by the Korea government Ministry of Science and ICT (NRF-2015M3C7A1028790, NRF-2018M3C7A1024603).

## CONFLICTS OF INTEREST

The authors declare no conflict of interest.

## FIGURE LEGENDS

**Fig. 1.** Neuronal functions of CYFIP2 and potential effects of the Arg87 variants identified in the West syndrome on those functions. (A) The Arg87 variants weaken the inhibitory CYFIP2-VCA interaction, thereby leading to aberrant activation of WRC and the downstream actin polymerization. This may result in dendritic spine remodeling and change excitatory/inhibitory synaptic balance, in favor of more excitation. (B) Similar to CYFIP1, CYFIP2 may also regulate neurotransmitter release in the presynaptic compartment, and the Arg87 variants may affect this function of CYFIP2. (C) The WRC interacts with certain ion channels potentially involved in regulating neuronal excitability. Given that CYFIP and ABI of the WRC form an interaction surface toward those membrane proteins, the Arg87 variants may indirectly affect such interactions, thereby changing surface stability or other functional properties of ion channels. (D) In addition to the WRC, CYFIP2 also interacts with some mRNA-binding proteins, such as FMRP, to regulate target mRNA transport and local translation. The Arg87 variants may affect formation of these mRNA-regulatory protein complexes, or their function in regulating target mRNAs.

## REFERENCES

1. Marin O, Valiente M, Ge X and Tsai LH (2010) Guiding neuronal cell migrations. Cold Spring Harbor perspectives in biology 2, a001834
2. Kevenaar JT and Hoogenraad CC (2015) The axonal cytoskeleton: from organization to function. Frontiers in molecular neuroscience 8, 44
3. Spence EF and Soderling SH (2015) Actin Out: Regulation of the Synaptic Cytoskeleton. The Journal of biological chemistry 290, 28613-28622
4. Yan Z, Kim E, Datta D, Lewis DA and Soderling SH (2016) Synaptic Actin Dysregulation, a Convergent Mechanism of Mental Disorders? The Journal of neuroscience : the official journal of the Society for Neuroscience 36, 11411-11417
5. Choi S-Y and Han K (2015) Emerging role of synaptic actin-regulatory pathway in the pathophysiology of mood disorders. Animal cells and systems 19, 283-288
6. Penzes P, Cahill ME, Jones KA, VanLeeuwen JE and Woolfrey KM (2011) Dendritic spine pathology in neuropsychiatric disorders. Nature neuroscience 14, 285-293
7. Takenawa T and Suetsugu S (2007) The WASP-WAVE protein network: connecting the membrane to the cytoskeleton. Nature reviews. Molecular cell biology 8, 37-48
8. Chen Z, Borek D, Padrick SB et al. (2010) Structure and control of the actin regulatory WAVE complex. Nature 468, 533-538
9. Mendoza MC (2013) Phosphoregulation of the WAVE regulatory complex and signal integration. Seminars in cell & developmental biology 24, 272-279
10. Krause M and Gautreau A (2014) Steering cell migration: lamellipodium dynamics and the regulation of directional persistence. Nature reviews. Molecular

cell biology 15, 577-590

11. Lee Y, Kim D, Ryu JR et al. (2017) Phosphorylation of CYFIP2, a component of the WAVE-regulatory complex, regulates dendritic spine density and neurite outgrowth in cultured hippocampal neurons potentially by affecting the complex assembly. *Neuroreport* 28, 749-754
12. Schenck A, Bardoni B, Moro A, Bagni C and Mandel JL (2001) A highly conserved protein family interacting with the fragile X mental retardation protein (FMRP) and displaying selective interactions with FMRP-related proteins FXR1P and FXR2P. *Proceedings of the National Academy of Sciences of the United States of America* 98, 8844-8849
13. Bagni C, Tassone F, Neri G and Hagerman R (2012) Fragile X syndrome: causes, diagnosis, mechanisms, and therapeutics. *The Journal of clinical investigation* 122, 4314-4322
14. Napoli I, Mercaldo V, Boyl PP et al. (2008) The fragile X syndrome protein represses activity-dependent translation through CYFIP1, a new 4E-BP. *Cell* 134, 1042-1054
15. De Rubeis S, Pasciuto E, Li KW et al. (2013) CYFIP1 coordinates mRNA translation and cytoskeleton remodeling to ensure proper dendritic spine formation. *Neuron* 79, 1169-1182
16. Abekhoukh S and Bardoni B (2014) CYFIP family proteins between autism and intellectual disability: links with Fragile X syndrome. *Frontiers in cellular neuroscience* 8, 81
17. Yoon KJ, Nguyen HN, Ursini G et al. (2014) Modeling a genetic risk for schizophrenia in iPSCs and mice reveals neural stem cell deficits associated with

- adherens junctions and polarity. *Cell stem cell* 15, 79-91
18. Oguro-Ando A, Rosensweig C, Herman E et al. (2015) Increased CYFIP1 dosage alters cellular and dendritic morphology and dysregulates mTOR. *Molecular psychiatry* 20, 1069-1078
  19. Bozdagi O, Sakurai T, Dorr N, Pilorge M, Takahashi N and Buxbaum JD (2012) Haploinsufficiency of *Cyfp1* produces fragile X-like phenotypes in mice. *PloS one* 7, e42422
  20. Chung L, Wang X, Zhu L et al. (2015) Parental origin impairment of synaptic functions and behaviors in cytoplasmic FMRP interacting protein 1 (*Cyfp1*) deficient mice. *Brain research* 1629, 340-350
  21. Kumar V, Kim K, Joseph C et al. (2013) C57BL/6N mutation in Cytoplasmic FMRP interacting protein 2 regulates cocaine response. *Science* 342, 1508-1512
  22. Han K, Chen H, Gennarino VA, Richman R, Lu HC and Zoghbi HY (2015) Fragile X-like behaviors and abnormal cortical dendritic spines in Cytoplasmic FMR1-interacting protein 2-mutant mice. *Human molecular genetics* 24, 1813-1823
  23. Zhang Y, Kang H, Lee Y et al. (2019) Smaller body size, early postnatal lethality, and cortical extracellular matrix-related gene expression changes of *Cyfp2*-null embryonic mice. *Frontiers in molecular neuroscience* in press
  24. Nakashima M, Kato M, Aoto K et al. (2018) De Novo Hotspot Variants in CYFIP2 Cause Early-Onset Epileptic Encephalopathy. *Annals of neurology*
  25. Peng J, Wang Y, He F et al. (2018) Novel West syndrome candidate genes in a Chinese cohort. *CNS neuroscience & therapeutics*
  26. Pathania M, Davenport EC, Muir J, Sheehan DF, Lopez-Domenech G and Kittler JT (2014) The autism and schizophrenia associated gene CYFIP1 is critical for

the maintenance of dendritic complexity and the stabilization of mature spines.

Translational psychiatry 4, e374

27. Sala C and Segal M (2014) Dendritic spines: the locus of structural and functional plasticity. *Physiological reviews* 94, 141-188
28. Collins MO, Husi H, Yu L et al. (2006) Molecular characterization and comparison of the components and multiprotein complexes in the postsynaptic proteome. *Journal of neurochemistry* 97 Suppl 1, 16-23
29. Bayes A, van de Lagemaat LN, Collins MO et al. (2011) Characterization of the proteome, diseases and evolution of the human postsynaptic density. *Nature neuroscience* 14, 19-21
30. Li J, Zhang W, Yang H et al. (2017) Spatiotemporal profile of postsynaptic interactomes integrates components of complex brain disorders. *Nature neuroscience* 20, 1150-1161
31. Han K, Holder JL, Jr., Schaaf CP et al. (2013) SHANK3 overexpression causes manic-like behaviour with unique pharmacogenetic properties. *Nature* 503, 72-77
32. Huttlin EL, Jedrychowski MP, Elias JE et al. (2010) A tissue-specific atlas of mouse protein phosphorylation and expression. *Cell* 143, 1174-1189
33. Schenck A, Bardoni B, Langmann C, Harden N, Mandel JL and Giangrande A (2003) CYFIP/Sra-1 controls neuronal connectivity in *Drosophila* and links the Rac1 GTPase pathway to the fragile X protein. *Neuron* 38, 887-898
34. Pan L, Zhang YQ, Woodruff E and Broadie K (2004) The *Drosophila* fragile X gene negatively regulates neuronal elaboration and synaptic differentiation. *Current biology : CB* 14, 1863-1870
35. Abekhoukh S, Sahin HB, Grossi M et al. (2017) New insights into the regulatory

- function of CYFIP1 in the context of WAVE- and FMRP-containing complexes. *Disease models & mechanisms* 10, 463-474
36. Zhao L, Wang D, Wang Q, Rodal AA and Zhang YQ (2013) *Drosophila* cyfip regulates synaptic development and endocytosis by suppressing filamentous actin assembly. *PLoS genetics* 9, e1003450
  37. Trowe T, Klostermann S, Baier H et al. (1996) Mutations disrupting the ordering and topographic mapping of axons in the retinotectal projection of the zebrafish, *Danio rerio*. *Development* 123, 439-450
  38. Baier H, Klostermann S, Trowe T, Karlstrom RO, Nusslein-Volhard C and Bonhoeffer F (1996) Genetic dissection of the retinotectal projection. *Development* 123, 415-425
  39. Pittman AJ, Gaynes JA and Chien CB (2010) *nev* (*cyfip2*) is required for retinal lamination and axon guidance in the zebrafish retinotectal system. *Developmental biology* 344, 784-794
  40. Cioni JM, Wong HH, Bressan D, Kodama L, Harris WA and Holt CE (2018) Axon-Axon Interactions Regulate Topographic Optic Tract Sorting via CYFIP2-Dependent WAVE Complex Function. *Neuron* 97, 1078-1093 e1076
  41. Marsden KC, Jain RA, Wolman MA et al. (2018) A *Cyfip2*-Dependent Excitatory Interneuron Pathway Establishes the Innate Startle Threshold. *Cell reports* 23, 878-887
  42. Kirkpatrick SL, Goldberg LR, Yazdani N et al. (2017) Cytoplasmic FMR1-Interacting Protein 2 Is a Major Genetic Factor Underlying Binge Eating. *Biological psychiatry* 81, 757-769
  43. Keane TM, Goodstadt L, Danecek P et al. (2011) Mouse genomic variation and

- its effect on phenotypes and gene regulation. *Nature* 477, 289-294
44. Chen B, Chou HT, Brautigam CA et al. (2017) Rac1 GTPase activates the WAVE regulatory complex through two distinct binding sites. *eLife* 6
  45. Darnell JC, Van Driesche SJ, Zhang C et al. (2011) FMRP stalls ribosomal translocation on mRNAs linked to synaptic function and autism. *Cell* 146, 247-261
  46. Lee JH, Kim HJ, Yoon JM et al. (2016) Interstitial deletion of 5q33.3q35.1 in a boy with severe mental retardation. *Korean journal of pediatrics* 59, S19-S24
  47. Spranger S, Rommel B, Jauch A, Bodammer R, Mehl B and Bullerdiek J (2000) Interstitial deletion of 5q33.3q35.1 in a girl with mild mental retardation. *American journal of medical genetics* 93, 107-109
  48. Focking M, Lopez LM, English JA et al. (2015) Proteomic and genomic evidence implicates the postsynaptic density in schizophrenia. *Molecular psychiatry* 20, 424-432
  49. Tiwari SS, Mizuno K, Ghosh A et al. (2016) Alzheimer-related decrease in CYFIP2 links amyloid production to tau hyperphosphorylation and memory loss. *Brain : a journal of neurology* 139, 2751-2765
  50. Gursoy S and Ercal D (2016) Diagnostic Approach to Genetic Causes of Early-Onset Epileptic Encephalopathy. *Journal of child neurology* 31, 523-532
  51. McTague A, Howell KB, Cross JH, Kurian MA and Scheffer IE (2016) The genetic landscape of the epileptic encephalopathies of infancy and childhood. *The Lancet. Neurology* 15, 304-316
  52. Bonaccorso CM, Spatuzza M, Di Marco B et al. (2015) Fragile X mental retardation protein (FMRP) interacting proteins exhibit different expression

- patterns during development. International journal of developmental neuroscience : the official journal of the International Society for Developmental Neuroscience 42, 15-23
53. Zhang Y, Chen K, Sloan SA et al. (2014) An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. The Journal of neuroscience : the official journal of the Society for Neuroscience 34, 11929-11947
  54. Bassell GJ and Warren ST (2008) Fragile X syndrome: loss of local mRNA regulation alters synaptic development and function. Neuron 60, 201-214
  55. Cajigas IJ, Tushev G, Will TJ, tom Dieck S, Fuerst N and Schuman EM (2012) The local transcriptome in the synaptic neuropil revealed by deep sequencing and high-resolution imaging. Neuron 74, 453-466
  56. Nelson SB and Valakh V (2015) Excitatory/Inhibitory Balance and Circuit Homeostasis in Autism Spectrum Disorders. Neuron 87, 684-698
  57. Lee Y, Zhang Y, Kim S and Han K (2018) Excitatory and inhibitory synaptic dysfunction in mania: an emerging hypothesis from animal model studies. Experimental & molecular medicine 50, 12
  58. Hsiao K, Harony-Nicolas H, Buxbaum JD, Bozdagi-Gunal O and Benson DL (2016) Cyfip1 Regulates Presynaptic Activity during Development. The Journal of neuroscience : the official journal of the Society for Neuroscience 36, 1564-1576
  59. Chen B, Brinkmann K, Chen Z et al. (2014) The WAVE regulatory complex links diverse receptors to the actin cytoskeleton. Cell 156, 195-207
  60. Contractor A, Klyachko VA and Portera-Cailliau C (2015) Altered Neuronal and



Circuit Excitability in Fragile X Syndrome. Neuron 87, 699-715






Fig. 1.