

BMB Reports – Manuscript Submission

Manuscript Draft

**Manuscript Number:** BMB-20-231

**Title:** Ethanol inhibits Kv7.2/7.3 channel open probability by reducing the PI(4,5)P2 sensitivity of Kv7.2 subunit

**Article Type:** Article

**Keywords:** Kv7.2/7.3 channel; ethanol; tetraethylammonium; open probability; PI(4,5)P2

**Corresponding Author:** Byung-Chang Suh

**Authors:** Kwon-Woo Kim<sup>1</sup>, Byung-Chang Suh<sup>1,\*</sup>

**Institution:** <sup>1</sup>Department of Brain and Cognitive Sciences, Daegu Gyeongbuk Institute of Science and Technology (DGIST),

**Manuscript Type:** Article

**Title:** Ethanol inhibits Kv7.2/7.3 channel open probability by reducing the PI(4,5)P<sub>2</sub> sensitivity of Kv7.2 subunit

**Author's name:** Kwon-Woo Kim and Byung-Chang Suh\*

**Affiliation:** Department of Brain and Cognitive Sciences, Daegu Gyeongbuk Institute of Science and Technology (DGIST), Daegu 42988, Korea

**Running Title:** Regulation of Kv7.2/7.3 channel properties by EtOH

**Keywords:** Kv7.2/7.3 channel; ethanol; tetraethylammonium; open probability; PI(4,5)P<sub>2</sub>.

**Corresponding Author's Information:** +82-53-785-6123 and bcsuh@dgist.ac.kr

**ABSTRACT**

Ethanol often causes critical health problems by altering the neuronal activities of the central and peripheral nerve systems. One of the cellular targets of ethanol is the plasma membrane proteins including ion channels and receptors. Recently, we reported that ethanol elevates membrane excitability in sympathetic neurons by inhibiting Kv7.2/7.3 channels in a cell type-specific manner. Even though our studies revealed that the inhibitory effects of ethanol on the Kv7.2/7.3 channel was diminished by the increase of plasma membrane phosphatidylinositol 4,5-bisphosphate (PI(4,5)P<sub>2</sub>), the molecular mechanism of ethanol on Kv7.2/7.3 channel inhibition remains unclear. By investigating the kinetics of Kv7.2/7.3 current in high K<sup>+</sup> solution, we found that ethanol inhibited Kv7.2/7.3 channels through a mechanism distinct from that of tetraethylammonium (TEA) which enters into the pore and blocks the gate of the channels. Using a non-stationary noise analysis (NSNA), we demonstrated that the inhibitory effect of ethanol is the result of reduction of open probability ( $P_o$ ) of the Kv7.2/7.3 channel, but not of a single channel current ( $i$ ) or channel number ( $N$ ). Finally, ethanol selectively facilitated the kinetics of Kv7.2 current suppression by voltage-sensing phosphatase (VSP)-induced PI(4,5)P<sub>2</sub> depletion, while it slowed down Kv7.2 current recovery from the VSP-induced inhibition. Together our results suggest that ethanol regulates neuronal activity through the reduction of open probability and PI(4,5)P<sub>2</sub> sensitivity of Kv7.2/7.3 channels.

## INTRODUCTION

Ethanol uptake acutely and chronically changes behavioral and psychological activities, such as mild behavior disinhibition, sedation, amnesia, and unconsciousness (1–3). Early studies focused on the effects of ethanol on the perturbation of membrane lipids in producing the physiological results at high concentrations (4). Recently, it has been revealed that ethanol regulates the activity of specific membrane proteins, such as receptors and voltage-gated and ligand-gated ion channels (4, 5). Ethanol treatment on the ventral tegmental neurons increased spontaneous firing frequency through the inhibition of Kv7 channels (6). On the contrary, ethanol enhances the current of G protein-gated inwardly rectifying potassium (GIRK) channels by directly binding the hydrophobic alcohol-binding pocket (7).

The heterotetrameric Kv7.2/7.3 channel is a voltage-gated potassium channel widely expressed in the central and peripheral nervous systems. It was originally termed M-channel since the currents were fully inhibited by the activation of M<sub>1</sub> muscarinic receptor (M<sub>1</sub>R) (8). In neurons, the Kv7.2 and 7.3 channel subunits highly co-localize at axon initial segments (AIS) regions (9) and form a stable Kv7.2/7.3 channel to generate a slowly activating and non-inactivating outward current with subthreshold membrane voltage near –60 mV (10). Hence, the modulation of Kv7.2/7.3 channel gating plays a crucial role in controlling the membrane potential and neuronal excitability (9). Interestingly, it has been revealed that plasma membrane phosphatidylinositol 4,5-bisphosphate (PI(4,5)P<sub>2</sub>) is absolutely required for activating Kv7 channels via enhancing the open probability (11). The Kv7.2/7.3 current is thus almost fully inhibited by depletion of plasma membrane PI(4,5)P<sub>2</sub> through the activation of M<sub>1</sub>R or exogenously expressed voltage-sensing phosphatase (VSP) (12, 13). Several recent papers suggest that PI(4,5)P<sub>2</sub> controls the Kv7.2/7.3 channel by interacting with S2-S3 linker, S4-S5 linker, the proximal C-terminus, and the A-B helix linker on C-terminus (14, 15).

We recently reported that ethanol reduces the outward K<sup>+</sup> current through Kv7.2/7.3

channels and elevates the membrane excitability in superior cervical ganglion (SCG) neurons, which is dependent upon the expression amount of Kv7 channels (16). Although we showed that ethanol inhibition of Kv7.2/7.3 channels was antagonized by PI(4,5)P<sub>2</sub> elevation in a concentration-dependent manner, underlying mechanisms of ethanol regulation remains unclear. In the present study, we revealed that ethanol suppresses the open probability of Kv7.2/7.3 channels through regulating gating activity without binding the channel pores directly. We also provided evidence that ethanol reduces the PI(4,5)P<sub>2</sub> sensitivity of Kv7.2/7.3 channels with the inhibitory order of Kv7.2 > Kv7.2/7.3 > Kv7.3 channels. These findings enhance our understanding of the molecular mechanisms underlying the ethanol-induced inhibitory modulation of Kv7 channel gating and neuronal excitability.

## RESULTS

### **Ethanol inhibition of Kv7.2/7.3 channels differs from block with internal TEA**

We examined whether the inhibitory action of ethanol is due to the direct blocking Kv7.2/7.3 channel pores as with TEA. TEA binds reversibly and independently to the outer and inner mouths of the pore and inhibit the currents through K<sup>+</sup> channels (17, 18). By increasing extracellular K<sup>+</sup> concentration, we first confirmed that changing the direction of the current does not alter the inhibitory mechanism by extracellular ethanol and TEA (see [Supplementary text and Fig. S1](#)). Since ethanol inhibits Kv7 current dose-dependently from 30 mM to 400 mM in both primary SCG neurons and tsA201 cell line (16), [here](#) we applied 400 mM of ethanol to tsA201 cells for studying the inhibitory mechanism of ethanol.

As cells were dialyzed internally with TEA, there were changes both in current waveform and in rectification (19). When TEA is drawn into the inner mouth of the pore at positive potentials, it blocks outward flow of K<sup>+</sup>. In contrast, when TEA is expelled from the inner mouth back into the cytoplasm at negative potentials, it allows inward flow of K<sup>+</sup> through

the pores but generates a “hook” shape current because of the ratification. Therefore, we tested for any current modification due to pore blockage by ethanol. For this, cells were dialyzed internally with 400 mM ethanol in the high-K<sup>+</sup> solution. However, there was no difference in the kinetics of Kv7.2/7.3 traces (Fig. 1A and 1C). On the other hand, with 1 mM internal TEA, the kinetics of the Kv7.2/7.3 current were significantly changed. (Fig. 1B and 1C). For examining the action of intracellular ethanol, Fig. 1D shows that after 3–5 min of TEA dialysis, the inward tail current had developed a hook at –70 mV. However, we found that intercellular ethanol did not change the current waveforms at any voltage application, indicating that the inhibition mechanism of ethanol is not due to direct binding to and blocking the pore of channel.

#### **Ethanol reduces the open probability of the Kv7.2/7.3 channel**

To further understand the mechanism of ethanol inhibition of Kv7.2/7.3 channels, we investigated single channel properties using the non-stationary noise analysis (NSNA) in the absence and presence of ethanol. As shown in Fig. 2A and 2B, the saturated Kv7.2/7.3 current was inhibited by  $37.6 \pm 1.9\%$  with ethanol treatment in normal Ringer’s solution. For performing the NSNA, the Kv7.2/7.3 current was measured at –20 mV for 2 s without or with ethanol and its variance was analyzed under each condition (Fig. 2C). Then, the variance-open probability relationship was fitted to the noise parabola equation for estimation of the maximum open probability ( $P_{Omax}$ ), the single channel current amplitude ( $i$ ) and the channel numbers ( $N$ ) (Fig. 2D). Our data showed that ethanol did not alter the single channel current or the numbers of Kv7.2/7.3 channel (Fig. 2E and 2F). However, ethanol significantly inhibited the maximum open probability of Kv7.2/7.3 channel (Fig. 2G). Together, the non-stationary noise analysis showed that ethanol inhibited the Kv7.2/7.3 current by reducing the open probability of the Kv7.2/7.3 channel.

#### **Ethanol inhibition of the Kv7.2/7.3 current is the result of reduction in PI(4,5)P<sub>2</sub> binding**

### sensitivity of the Kv7.2/7.3 channel

We previously showed that the inhibitory effect of ethanol on the Kv7.2/7.3 current is modulated by the concentration of the plasma membrane PI(4,5)P<sub>2</sub> by activating *Ciona intestinalis* voltage-sensing phosphatase (*Ci*-VSP) (16). However, the acute effects of ethanol on the kinetics of PI(4,5)P<sub>2</sub> degradation and resynthesis in the plasma membrane was not clearly solved. Here, in order to further understand how ethanol alters the kinetics of Kv7.2/7.3 current inhibition by VSP activation, whole-cell patch clamp and FRET imaging were simultaneously performed in the same living cell and the changes of Kv7.2/7.3 current and PI(4,5)P<sub>2</sub> concentration during and after VSP activation were measured at the same time. The perturbation of plasma membrane PI(4,5)P<sub>2</sub> level was detected using Förster resonance energy transfer (FRET) assay between two PI(4,5)P<sub>2</sub> probes, CFP-PH(PLCδ1) and YFP-PH(PLCδ1), as described previously (20, 21). The results showed that Kv7.2/7.3 current inhibition upon activation of *Danio rerio* VSP (*Dr*-VSP) by depolarization to +50 mV for 40 s was facilitated approximately 2-fold by ethanol treatment (Fig. 3A-top and 3B-left). However, FRET imaging in response to PI(4,5)P<sub>2</sub> degradation to PI(4)P by VSP activation was not significantly changed (Fig. 3A-bottom and 3B-right). Similarly, ethanol slowed Kv7.2/7.3 recovery after VSP activation, whereas the PI(4,5)P<sub>2</sub> resynthesis indicated by FRET<sub>r</sub> was not changed (Fig. 3C and 3D), suggesting that ethanol decreases the PI(4,5)P<sub>2</sub> sensitivity of the Kv7 channel as the inhibitory effects of ethanol are stronger at low PI(4,5)P<sub>2</sub> concentration in the plasma membrane.

Ethanol inhibition of current differs in each subtype of Kv7 channel and is inversely correlated with their PI(4,5)P<sub>2</sub> binding affinity (16). By using this characteristic, we further examined if ethanol differentially regulates the VSP-mediated current inhibition in Kv7 channel subtypes depending on their PI(4,5)P<sub>2</sub> binding affinity. As shown in Fig. 4, ethanol hastened the kinetics of VSP-induced current inhibition in Kv7.2 (Fig. 4A, and 4D), but not in

Kv7.3 channels (Fig. 4C and 4F). Heteromeric Kv7.2/7.3 channels had a significant reaction by ethanol (Fig. 4B and 4E). However, ethanol did not change the voltage-dependent activation of homomeric Kv7.2 and Kv7.3 channels (Supplementary Fig. S2), suggesting that ethanol selectively facilitated the inhibition of Kv7.2 channels with low apparent PI(4,5)P<sub>2</sub> binding affinity, but not Kv7.3 channels with high PI(4,5)P<sub>2</sub> binding affinity. Taken together, our results further indicated that ethanol inhibits Kv7.2/7.3 currents through the reduction of PI(4,5)P<sub>2</sub> sensitivity of the Kv7.2 subtype, which might be the molecular mechanism responsible for the ethanol-mediated K<sup>+</sup> current suppression through Kv7.2/7.3 channels.

## DISCUSSION

In this study, we demonstrated that ethanol inhibits Kv7.2/7.3 channels through a mechanism different from that of TEA. The kinetics analysis of current deactivation and activation in the presence of internal ethanol indicated that ethanol does not block the pore directly unlike TEA. We further confirmed that the ethanol inhibition of the Kv7.2/7.3 channel was caused by reduction of open probability of the Kv7.2/7.3 channel. We also obtained more definitive results showing that ethanol regulated the binding affinity between Kv7.2/7.3 channel and PI(4,5)P<sub>2</sub>. Our data further show how ethanol modulates the gating of the Kv7 channels in a subtype-specific and phospholipid-dependent manner.

In Kv7.2/7.3 channels, extracellular TEA inhibits currents by binding the tyrosine residue of 323 of the pore of Kv7.2, but not the Kv7.3 channel because it has a threonine at the corresponding position (17). Thus, the inhibitory effects of TEA are different on different Kv7 subtypes. The kinetics of internal TEA inhibition has been reported for the prokaryotic KcsA channel through the internal quaternary ammonium ions located at the internal water cavity and one of the ethyl groups inserted into the selectivity filter (22, 23). Moreover, the mechanism of ethanol regulation has been studied for several potassium channels including GIRK2 channel

and Kv3.4 channel. There are conserved hydrophobic alcohol-binding pockets exist in the cytoplasmic domain of GIRK2 channel (24, 25). Studies of the Kv3.4 channel indicate that the substitution of glycine at 371 of the S4-S5 linker to isoleucine alters the alcohol-sensitive channel (26). For these reasons, the ethanol binding sites of Kv7.2/7.3 channel **might not be** related to the pore region where TEA binds to block the current.

The properties of single channel appeared to be consistent with the literature under similar conditions of membrane potential (27). There are different results for single channel current of Kv7.2/7.3 channel, such as  $\sim 0.21$  pA at  $-33$  mV using NSNA or  $\sim 0.5$  pA using the single channel recording technique (11, 27, 28). According to our data, the single channel current and the open probability were calculated for  $\sim 0.4$  pA and 0.4 at  $-20$  mV of membrane potential, respectively. The diversity in single channel amplitude indicates a result of the heterogeneous Kv7.2 and Kv7.3 channel complex (29, 30). The correlation between the regulation of amplitude of Kv7.2/7.3 currents and open probability by several drugs has been investigated. The single channel patch clamp technique showed that the open probability of Kv7.2/7.3 channel was increased in the presence of the positive allosteric modulator retigabine, a water soluble PI(4,5)P<sub>2</sub> analog, DiC<sub>8</sub>-PI(4,5)P<sub>2</sub>, or intracellular zinc (11, 28, 31). According to our data, ethanol inhibited the open probability of the Kv7.2/7.3 channel rather than regulating single channel current or channel numbers.

Through simultaneous measurement of the Kv7 current and PI(4,5)P<sub>2</sub> changes using patch clamp and FRET imaging, we further confirmed that the facilitatory effects of ethanol are not due to the elevation of VSP activity or inhibition of enzymes involved in the PI(4,5)P<sub>2</sub> metabolism pathways in the cell membrane. Rather, our results suggest that the ethanol effects on the Kv channel under the VSP activation are the result of reducing the PI(4,5)P<sub>2</sub> binding affinity for the Kv7 channels. This was further confirmed with ethanol experiments with Kv7 subtypes. Usually Kv7.3 has a more than 20-fold higher apparent affinity for PI(4,5)P<sub>2</sub> than

Kv7.2 (32). Consistently, we found that the ethanol facilitation of K<sup>+</sup> current inhibition by VSP was much stronger in Kv7.2. The basic residues on the S2-S3 linker, the S4-S5 linker, and the C-terminus region of Kv7.2/7.3 channel have been suggested as the sites for the interaction with PI(4,5)P<sub>2</sub> by charge neutralization or deletion using a mutagenesis technique (11, 15, 32). However, it is still unclear how ethanol regulates the interaction of PI(4,5)P<sub>2</sub> with Kv7.2 subtypes.

In summary, we showed here that ethanol suppresses Kv7.2/7.3 channel activity by reducing the open probability and PI(4,5)P<sub>2</sub> sensitivity of Kv7.2/7.3 channels. Because ethanol affects acute and chronic physiological functions in excitable cells, including hormone secretion from glands, neural firing and neurotransmitter release, and vascular systems, determining the regulatory mechanism of Kv7 channel suppression by ethanol will provide an important clue to understanding such diverse effects of ethanol in physiology and pathophysiology.

## MATERIALS AND METHODS

Additional materials and methods are included in the supplementary materials.

### Electrophysiological recording

The whole cell patch clamp technique was performed at room temperature using an EPC-10 patch clamp amplifier (HEKA Elektronik, Germany). The electrodes were pulled from glass micropipette capillaries using a P-97 micropipette puller (Sutter Instrument, USA) with a resistance of 2–4 MΩ. Series-resistance errors were compensated by 60%. The average cell capacitance used in this study was  $17.2 \pm 0.65$  pF. The Kv7 current was measured at –20 mV and in a 500 ms hyperpolarizing step to –60 mV every 2 or 4 s, as described previously (16).

### Förster resonance energy transfer (FRET)

The FRET experiments were simultaneously performed while measuring the Kv7.2/7.3 current in a single cell. The FRET signals were acquired, and real time was calculated by a homemade program as previously described (33). The FRET calculated ratio (cFactor = CFP/YFP = 0.55) was used to adjust the raw YFP emission signal. The signal of the FRET ratio was calculated as follows:  $FRET_r = (YFP_c - cFactor \times CFP_c)/CFP_c$ , where YFP<sub>c</sub> is the signal from YFP excited from the FRET signal (YFP emission by CFP excitation), YFP emission is detected by the long wavelength photomultiplier (YFP<sub>c</sub>), and CFP emission is detected by the short wavelength photomultiplier.

### Non-stationary noise analysis

Non-stationary noise analysis (NSNA) was performed as described (34, 35). Kv7.2/7.3 channels were depolarized every 3 s with  $-20$  mV pulse during 2 s for channel activation from  $-80$  mV holding potential. Twenty to forty traces of Kv7 current were collected when fully stabilized in the absence or presence of ethanol. The Kv7 currents were low-pass filtered at 10 kHz and digitized with a sampling rate of 100 kHz for acquiring sufficient data. The mean current versus variance was binned into 30 bins, and data were fit with the following equation:  $\sigma^2 = iI - I^2/N + \sigma_b^2$  ( $\sigma^2$ : variance,  $i$ : single channel current,  $I$ : mean current,  $N$ : total number of channels,  $\sigma_b^2$ : background variance). The maximum open probability,  $P_{Omax}$ , of the peak current can be calculated from the following equation:  $P_{Omax} = I_{max}/(i \times N)$ , where  $I_{max}$  is the maximum value of mean current. For presenting plots of variance/ $\langle I \rangle_{max}$  versus open probability, the background variance ( $\sigma_b^2$ ) is subtracted from variance and each individual trace was divided by the estimated  $\langle I \rangle_{max}$  to represent the open probability and variance/ $\langle I \rangle_{max}$  (35).

### ACKNOWLEDGMENTS

This work was supported by grants from the National Research Foundation of Korea (NRF)

grant funded by the Korea government (MSIP) (No. 2019R1A2B5B01070546), the Basic Science Research Program (2020R1A4A1019436), the Korea Brain Research Institute (KBRI) basic research program funded by the Ministry of Science, ICT and Future Planning to JCR (No. 20-BR-04-01).

## CONFLICTS OF INTEREST

The authors declare no conflict of interest.

## FIGURE LEGENDS

**Fig. 1.** Intracellular TEA and EtOH show differential effects on the activation and deactivation kinetics of Kv7.2/7.3 current in high-K<sup>+</sup> external solution. Comparison of Kv7.2/7.3 current traces obtained from cells intracellularly perfused with vehicle (control) or 400 mM EtOH (A) and 1 mM TEA (B). (C) Summary of time constants for deactivation and activation in cells perfused with 400 mM EtOH or 1 mM TEA. (control,  $n = 8$ ; 400 mM EtOH,  $n = 8$ ; 1 mM TEA,  $n = 4$ ). One-way analysis of variance followed by Sidak's posthoc test, \*\*  $p < 0.01$ , as compared with control. n.s., non-significance. (D) Deactivation kinetics of inward current at  $-70$  mV after  $-20$  mV (left) or  $+40$  mV (right) pulse step in high-K<sup>+</sup> external solution. Dotted line indicates Kv7.2/7.3 current in control cells. Dashed line is the zero-current level.

**Figure 2.** Non-stationary noise analysis (NSNA) for whole cell configuration of Kv7.2/7.3 channel with or without EtOH treatment. (A) Left, time-dependent regulation of Kv7.2/7.3 current upon treatment of 400 mM EtOH for 150 s. Right, summary of current inhibition by 400 mM EtOH ( $n = 9$ ). (B) Representative traces of Kv7.2/7.3 current (upper panel) and variance (under panel) elicited in the absence (black) and presence (blue) of 400 mM EtOH in normal Ringer's solution. (C) The  $\text{variance}/\langle I \rangle_{\max}$  plotted against the open probability of

Kv7.2/7.3 channel without (left) or with (right) 400 mM EtOH. Mean single channel current ( $i$ ; E), channel number ( $N$ ; F), and maximal open probability ( $P_{o,max}$ ; G) were calculated by non-stationary noise analysis ( $n = 9$ ). Paired Student's  $t$ -test were performed. \*\*\*  $p < 0.001$ .

**Figure 3.** Simultaneous measurement of EtOH effects on the regulation of Kv7.2/7.3 current and membrane PIP<sub>2</sub> by *Danio rerio* voltage-sensing phosphatase (Dr-VSP) in a single cell. (A) Representative current traces (top) and FRETr (bottom) without (grey or black lines) or with (blue lines) 400 mM EtOH during PI(4,5)P<sub>2</sub> degradation by Dr-VSP. (B) Time constants ( $\tau$ ) for Kv7.2/7.3 current inhibition (left) and FRETr changes (right) caused by activation of Dr-VSP in the absence (grey bars) or presence (blue bars) of 400 mM EtOH. Current: control,  $n = 8$ ; ethanol,  $n = 7$ ; FRETr: control,  $n = 6$ ; ethanol,  $n = 6$ . (C) Representative current traces (top) and FRETr (bottom) measured at  $-20$  mV for 40 s after depolarization to  $+100$  mV for 1 s to deplete membrane PI(4,5)P<sub>2</sub> in cells expressing Dr-VSP. (D) Half recovery ( $1/2$ ) for Kv7.2/7.3 current (left) and FRETr changes (right) after activation of Dr-VSP in the absence (grey bars) and presence (blue bars) of 400 mM EtOH,  $n = 4$  each. Mann–Whitney  $u$ -test, \*  $p < 0.05$ , \*\*  $p < 0.01$ .

**Figure 4.** Kinetics of Kv7 subtype current inhibition facilitated by VSP activation. (A, B, C) Cells were cells transfected with *Ciona intestinalis* VSP (Ci-VSP) plus Kv7.2 (A), both Kv7.2 and Kv7.3 (B), or Kv7.3 alone (C) and depolarized to  $+30$  mV for 42 s to deplete PI(4,5)P<sub>2</sub> through Ci-VSP activation in the absence (black) or presence (blue) of EtOH. Relative current traces of the Kv7 channels subtracted to endogenous potassium current with Ci-VSP transfected are superimposed. (D, E, F) Time constants ( $\tau$ ) of Kv7 current inhibition by the activation of Ci-VSP in cells expressing Kv7.2 ( $n = 4$  each) (D), both Kv7.2 and Kv7.3 ( $n = 6$  for control, 7 for EtOH) (E), or Kv7.3 alone ( $n = 4$  each) (F) without (black bars) and with (blue bars) 400 mM EtOH. Mann–Whitney  $u$ -test, \*  $p < 0.05$ , \*\*  $p < 0.01$ .

## REFERENCES

1. Harrison NL, Skelly MJ, Grosserode EK et al (2017) Effects of acute alcohol on excitability in the CNS. *Neuropharmacology* 122, 36–45
2. Alderazi Y and Brett F (2007) Alcohol and the nervous system. *Curr Diagn Pathol* 13, 203–209
3. Hirota Y (1976) Effect of ethanol on contraction and relaxation of isolated rat ventricular muscle. *J Mol Cell Cardiol* 8, 727–732
4. Harris RA, Trudell JR and Mihic SJ (2008) Ethanol's molecular targets. *Sci Signal* 1, re7
5. Kruse SW, Zhao R, Smith DP and Jones DNM (2003) Structure of a specific alcohol-binding site defined by the odorant binding protein LUSH from *Drosophila melanogaster*. *Nat Struct Mol Biol* 10, 694–700
6. Bodhinathan K and Slesinger PA (2013) Molecular mechanism underlying ethanol activation of G-protein-gated inwardly rectifying potassium channels. *Proc Natl Acad Sci* 110, 18309–18314
7. Koyama S, Brodie MS and Appel SB (2007) Ethanol inhibition of M-current and ethanol-induced direct excitation of ventral tegmental area dopamine neurons. *J Neurophysiol* 97, 1977–1985
8. Brown DA and Adams PR (1980) Muscarinic suppression of a novel voltage-sensitive  $K^+$  current in a vertebrate neurone. *Nature*, 673–6.
9. Greene DL and Hoshi N (2017) Modulation of Kv7 channels and excitability in the brain. *Cell Mol Life Sci* 74, 495-508
10. Wang H-S, Pan Z, Shi W, Brown BS et al (1998) KCNQ2 and KCNQ3 potassium channel subunits: molecular correlates of the M-channel. *Science* 282, 1890–1893
11. Telezhkin V, Brown DA and Gibb AJ (2012) Distinct subunit contributions to the activation

- of M-type potassium channels by PI(4,5)P<sub>2</sub>. *J Gen Physiol* 140, 41–53
12. Suh BC and Hille B (2002) Recovery from muscarinic modulation of M current channels requires phosphatidylinositol 4,5-bisphosphate synthesis. *Neuron* 35, 507–520
13. Falkenburger BH, Jensen JB and Hille B (2010) Kinetics of PIP<sub>2</sub> metabolism and KCNQ2/3 channel regulation studied with a voltage-sensitive phosphatase in living cells. *J Gen Physiol* 135, 99–114
14. Zhang Q, Zhou P, Chen Z et al (2013) Dynamic PIP<sub>2</sub> interactions with voltage sensor elements contribute to KCNQ2 channel gating. *Proc Natl Acad Sci* 110, 20093–20098
15. Choveau FS, De la Rosa V, Bierbower SM, Hernandez CC and Shapiro MS (2018) Phosphatidylinositol 4,5-bisphosphate (PIP<sub>2</sub>) regulates KCNQ3 K<sup>+</sup> channels by interacting with four cytoplasmic channel domains. *J Biol Chem* 293, 19411–19428
16. Kim KW, Kim K, Lee H and Suh BC (2019) Ethanol elevates excitability of superior cervical ganglion neurons by inhibiting Kv7 channels in a cell type-specific and PI(4,5)P<sub>2</sub>-dependent manner. *Int J Mol Sci* 20, 4419
17. Hadley JK, Noda M, Selyanko AA et al (2000) Differential tetraethylammonium sensitivity of KCNQ1–4 potassium channels. *Br J Pharmacol* 129, 413–415
18. Holmgren M, Smith PL and Yellen G (1997) Trapping of organic blockers by closing of voltage-dependent K<sup>+</sup> channels: Evidence for a trap door mechanism of activation gating. *J Gen Physiol* 109, 527–535
19. Suh BC and Hille B (2007) Electrostatic Interaction of internal Mg<sup>2+</sup> with membrane PIP<sub>2</sub> seen with KCNQ K<sup>+</sup> channels. *J Gen Physiol* 130, 241–256
20. Keum D, Kruse M, Kim DI, Hille B and Suh BC (2016) Phosphoinositide 5- and 3-phosphatase activities of a voltage-sensing phosphatase in living cells show identical voltage dependence. *Proc Natl Acad Sci* 113, E3686–E3695
21. Jensen JB, Lyssand JS, Hague C and Hille B (2009) Fluorescence changes reveal kinetic

steps of muscarinic receptor-mediated modulation of phosphoinositides and Kv7.2/7.3 K<sup>+</sup> channels. *J Gen Physiol* 133, 347–359

22. Kutluay E, Roux B and Heginbotham L (2005) Rapid intracellular TEA block of the KcsA potassium channel. *Biophys J* 88, 1018–1029

23. Luzhkov VB and Åqvist J (2001) Mechanisms of tetraethylammonium ion block in the KcsA potassium channel. *FEBS Lett* 495, 191–196

24. Pegan S, Arrabit C, Slesinger PA and Choe S (2006) Andersen's syndrome mutation effects on the structure and assembly of the cytoplasmic domains of Kir2.1. *Biochemistry* 45, 8599–8606

25. Aryal P, Dvir H, Choe S and Slesinger PA (2009) A discrete alcohol pocket involved in GIRK channel activation. *Nat Neurosci* 12, 988–995

26. Covarrubias M, Vyas TB, Escobar L and Wei A (1995) Alcohols inhibit a cloned potassium channel at a discrete saturable site. Insights into the molecular basis of general anesthesia. *J Biol Chem* 270, 19408–19416

27. Villarroel A (1997) Nonstationary noise analysis of M currents simulated and recorded in PC12 cells. *J Neurophysiol* 77, 2131–2138

28. Tatulian L and Brown DA (2003) Effect of the KCNQ potassium channel opener retigabine on single KCNQ2/3 channels expressed in CHO cells. *J Physiol* 549, 57–63

29. Schwake M, Pusch M, Kharkovets T and Jentsch TJ (2000) Surface expression and single channel properties of KCNQ2/KCNQ3, M-type K<sup>+</sup> channels involved in epilepsy. *J Biol Chem* 275, 13343–13348

30. Selyanko AA, Hadley JK and Brown DA (2001) Properties of single M-type KCNQ2/KCNQ3 potassium channels expressed in mammalian cells. *J Physiol* 534, 15–24

31. Gao H, Boillat A, Huang D, Liang C, Peers C and Gamper N (2017) Intracellular zinc activates KCNQ channels by reducing their dependence on phosphatidylinositol 4,5-

bisphosphate. *Proc Natl Acad Sci* 114, E6410–E6419


32. Zhang H, Craciun LC, Mirshahi T et al (2003) PIP<sub>2</sub> activates KCNQ channels, and its hydrolysis underlies receptor-mediated inhibition of M currents. *Neuron* 37, 963–975


33. Keum D, Baek C, Kim DI, Kweon HJ and Suh BC (2014) Voltage-dependent regulation of Ca<sub>v</sub>2.2 channels by G<sub>q</sub>-coupled receptor is facilitated by membrane-localized β subunit. *J Gen Physiol* 144, 297–309


34. Alvarez O, Gonzalez C and Latorre R (2002) Counting channels: A tutorial guide on ion channel fluctuation analysis. *Adv Physio Educ* 26, 327–341


35. Hartveit E and Veruki ML (2007) Studying properties of neurotransmitter receptors by non-stationary noise analysis of spontaneous postsynaptic currents and agonist-evoked responses in outside-out patches. *Nat Protoc* 2, 434–448

36. Lim NK, Lam AKM and Dutzler R (2016) Independent activation of ion conduction pores in the double-barreled calcium-activated chloride channel TMEM16A. *J Gen Physiol.* 148, 375–392









**Ethanol inhibits Kv7.2/7.3 channel open probability by reducing the PI(4,5)P<sub>2</sub> sensitivity of Kv7.2 subunit**

Kwon-Woo Kim and Byung-Chang Suh\*

Department of Brain and Cognitive Sciences, Daegu Gyeongbuk Institute of Science and Technology (DGIST), Daegu 42988, Korea

\*Corresponding Author: +82-53-785-6123 and bcsuh@dgist.ac.kr

## Supplementary materials and methods

### MATERIALS AND METHODS

#### Cell Culture and Transfection

TsA201 cells were cultured in Dulbecco's Modified Eagle Medium (Invitrogen, Life Technologies, CA, USA) with 10% fetal bovine serum (HyClone, USA) and 0.2% penicillin/streptomycin (Invitrogen, Life Technologies, CA, USA) at 37°C under 5% CO<sub>2</sub>. Cells were transfected using Lipofectamine 2000 (Invitrogen, Life Technologies, CA, USA) with various DNAs, including human Kv7.2, human Kv7.3 channel subunits (1:1 ratio) and Ds-Red for a transfection marker. In some experiments, Ci-VSP or Dr-VSP with EFGP were co-transfected. For Förster resonance energy transfer (FRET) experiments, eYFP-PH(PLC $\delta$ 1) and CFP-PH(PLC $\delta$ 1) were used with Dr-VSP. The transfected cells were plated on poly L-lysine (Sigma-Aldrich, USA)-coated chips the day after transfection using trypsin. Plated cells were used for current recording experiments within 36–48 h after transfection, as described previously (16).

#### Solutions and Materials

The Ringer's solution was used to record the K<sup>+</sup> current and for FRET signal. It consisted of (in mM) 160 NaCl, 2.5 KCl, 2 CaCl<sub>2</sub>, 1 MgCl<sub>2</sub>, 10 HEPES, and 8 glucose, adjusted to pH 7.4 with NaOH. The pipette solution contained (in mM) 175 KCl, 5 MgCl<sub>2</sub>, 5 HEPES, 0.1 BAPTA, 3 Na<sub>2</sub>ATP, and 0.1 Na<sub>3</sub>GTP, then adjusted to pH 7.4 with KOH. The high-K<sup>+</sup> Ringer's solution contained (in mM) 132.5 NaCl, 30 KCl, 2 CaCl<sub>2</sub>, 1 MgCl<sub>2</sub>, 10 HEPES, and 8 glucose, adjusted to pH 7.4 with NaOH.


#### Data analysis

Data were analyzed with Excel 2016 and Igor Pro (WaveMetrics, Inc., USA). The time constants were measured by exponential fit. All quantitative data are expressed as mean  $\pm$  standard error. One-way analysis of variance (ANOVA) followed by Sidak's posthoc test and Mann–Whitney u-test, and paired Student's *t*-test were performed. Data are expressed as mean  $\pm$  SEM. The differences were considered significant at the  $p < 0.05$  level.


**Text for Supplementary Fig. S1**

Tetraethylammonium binds reversibly and independently to the outer and inner mouths of the pore and inhibit the currents through  $K^+$  channels (1, 2). Using high- $K^+$  Ringer's solution, we investigate the inhibitory mechanism of ethanol in Kv7 channels as compared to TEA. The extracellular  $K^+$  concentration was elevated to 30 mM to raise the  $K^+$  equilibrium potential to  $-45$  mV, so we could observe inward currents negative to  $-45$  mV. Thus, the tails at  $-70$  mV were changed from outward to inward current in 30 mM  $K^+$  external solution. However, it did not alter the voltage dependence of activation (Supplementary Fig. S1A-right) or time constants of activation and deactivation of the Kv7 current (Supplementary Fig. S1B). In addition, the ethanol inhibition of the Kv7.2/7.3 current was not changed in the extracellular high- $K^+$  concentration. The inward and outward currents were symmetrically inhibited in the presence of 200 mM and 400 mM ethanol, respectively (Supplementary Fig. S1C). The application of 30 mM TEA to the extracellular solution also symmetrically reduced the Kv7.2/7.3 current without any changes in the rates of activation or deactivation of the current (Supplementary Fig. S1D, also see inset). These results indicate that changing the direction of the current by increasing extracellular  $K^+$  concentration does not alter the inhibitory mechanism by extracellular ethanol and TEA.

1. Hadley JK, Noda M, Selyanko AA, Wood IC, Abogadie FC and Brown DA (2000) Differential tetraethylammonium sensitivity of KCNQ1–4 potassium channels. *Br J Pharmacol* 129, 413–415
2. Holmgren M, Smith PL and Yellen G (1997) Trapping of organic blockers by closing of voltage-dependent  $K^+$  channels: evidence for a trap door mechanism of activation gating. *J Gen Physiol* 109, 527–535



**Supplementary Fig. S1.** Symmetrical inhibition of Kv7.2/7.3 current in tsA201 cells by ethanol (EtOH) and tetraethylammonium (TEA) in high- $K^+$  external solution. (A) Left, families of Kv7.2/7.3 current in 2.6 (normal) and 30 mM (high)  $K^+$  Ringer's solution elicited by voltage steps from  $-70$  to  $+50$  mV, in 10-mV intervals. Inset shows the pulse protocol. The holding potential is  $-70$  mV. Right, relative tail currents measured between 10 and 20 ms after depolarization in 2.6 (normal) and 30 mM (high)  $K^+$  Ringer's solution,  $n = 4$  each. (B) Left, representative traces of Kv7.2/7.3 currents elicited by the protocol of  $+20$  to  $-60$  mV in normal (black) and high- $K^+$  (blue) solution. Right panels are their mean activation and deactivation time constants,  $n = 6$ . (C) Left, Symmetrical Kv7.2/7.3 current regulation by 200 mM and 400 mM EtOH treatments in 30 mM  $K^+$  Ringer's solution. Right, summary of ethanol inhibition of Kv7.2/7.3 current at  $-20$  mV and  $-60$  mV.  $n = 9$ . (D) Left, symmetrical Kv7.2/7.3 current inhibition by 30 mM TEA in high- $K^+$  Ringer's solution. Right, summary of Kv7.2/7.3 inhibition at  $-20$  mV and  $-60$  mV by TEA,  $n = 7$ . Data are expressed as mean  $\pm$  SEM.



**Supplementary Fig. S2.** Ethanol does not change the voltage-dependent activation of Kv7 channels. (A, D and G) Representative current traces in the absence (black) or presence (blue) of 400 mM ethanol in tsA201 cells transfected with Kv7.2 and Kv7.3 channel subunits either individually or in combination. Normalized conductance versus voltage relationship of Kv7.2 (B: n= 4, 5), Kv7.2/7.3 (E: n=11), and Kv7.3 (H: n=7). The Boltzmann functions was performed to fit the conductance-voltage curve and to determine the half-maximal activation voltage ( $V_{\text{half}}$ ) of Kv7 channel (C, F, and I).