

BMB Reports – Manuscript Submission

Manuscript Draft

Manuscript Number: BMB-16-078

Title: The novel function of Stabilin-2 in myoblast fusion: The recognition of extracellular phosphatidylserine as a “fuse-me” signal

Article Type: Perspective (Invited Only)

Keywords: Stabilin-2; Muscle regeneration; Stab2; Myoblast fusion; phosphatidylserine

Corresponding Author: In-San Kim

Authors: Go-Woon Kim¹, Seung-Yoon Park^{1,2}, In-San Kim^{1,3,*}

Institution: ¹Center for Theragnosis, Biomedical Research Institute, Korea Institute of Science and Technology (KIST), Seoul 02792, Republic of Korea,

²Department of Biochemistry, School of Medicine, Dongguk University, Gyeongju 780-714, Republic of Korea,

³KU-KIST school, Korea University, Seoul 02841, Republic of Korea,

The novel function of Stabilin-2 in myoblast fusion: The recognition of extracellular phosphatidylserine as a “fuse-me” signal

Go-Woon Kim¹, Seung-Yoon Park^{1,2} and In-San Kim^{1,3,*}

¹Center for Theragnosis, Biomedical Research Institute, Korea Institute of Science and Technology (KIST), Seoul 02792, Republic of Korea; ²Department of Biochemistry, School of Medicine, Dongguk University, Gyeongju 780-714, Republic of Korea; ³KU-KIST school, Korea University, Seoul 02841, Republic of Korea.

* Corresponding author. E-mail: iskim14@kist.re.kr.

Abstract

Myoblast fusion is important for skeletal muscle formation. Even though the knowledge of myoblast fusion mechanism has accumulated over the years, the initial signal of fusion is yet to be elucidated. Our study reveals the novel function of a phosphatidylserine (PS) receptor, stabilin-2 (Stab2) in the modulation of myoblast fusion through the recognition of PS exposed on myoblasts. During differentiation of myoblasts, Stab2 expression is higher than other PS receptors and is controlled by calcineurin/NFAT signaling on myoblasts. The forced expression of Stab2 results in an increase in myoblast fusion, and genetic ablation of Stab2 in mice causes a reduction in muscle size, as a result of impaired myoblast fusion. After muscle injury, muscle regeneration is impaired in Stab2-deficient mice, resulting in small myofibers with fewer nuclei, which is due to reduction of fusion rather than defection of myoblast differentiation. The fusion-promoting role of Stab2 is dependent on its PS-binding motif, and the blocking of PS-Stab2 binding impairs cell-cell fusion on myoblasts. Given our previous finding that Stab2 recognizes PS exposed on apoptotic cells for sensing as an “eat-me” signal, we propose that PS-Stab2 binding is for sensing of a “fuse-me” signal as the initial signal of myoblast fusion.

Skeletal muscle is formed from myogenic precursors, and after myogenic commitment, mono-nucleated myoblasts fuse with each other and form multinucleated myofibers. For the past two decades, numerous molecules have been suggested for the regulators of myoblast fusion, including membrane receptors, transcription factors, and signaling molecules. However, the details of fusion mechanism has been elusive.

Phosphatidylserine (PS) is one of the phospholipid membrane components and is normally localized at the inner leaflet of the plasma membrane. Exposure of PS to the outer plasma membrane has important roles in various cellular events, such as an “eat-me” signal for apoptotic cell clearance, a scaffold for coagulation factors in activated platelets, and a candidate signal for cell-cell fusion. In the case of myoblast fusion, the function of PS exposure has not much been analyzed but one thing is known, that transient exposure of PS on the extracellular membrane of fusing myoblasts is needed for myoblast fusion. A recent study showed that a PS-receptor, brain-specific angiogenesis inhibitor 1 (Bai1), recognized PS exposed on apoptotic cells and promoted fusion between healthy myoblasts which may result in recruitment of ELMO/Rac1 (Hochreiter-Hufford et al. (2013) *Nature* 497:263-267, doi:10.1038/nature12135). However, the receptor recognizing PS on healthy myoblasts has not been identified.

Our study has reported for the first time that the function of Stab2 in myoblast fusion is related to the recognition of PS exposed in healthy myoblasts. The expression level of Stab2 was higher than other PS receptors on mouse skeletal muscles and increased depending on the differentiation days of C2C12 and primary myoblasts. Stab2 expression was regulated by calcineurin/NFAT signaling; in detail, NFATc1 binding to the Stab2 promoter region enhanced the expression of Stab2 transcript. The forced expression of Stab2 increased the

fusion of C2C12 cells, but there were no differences in the expression of myogenin, one of the differentiation makers for committed myoblasts. Furthermore, mouse fibroblasts L cells, which usually does not fuse with each other, acquired fusogenic activity after the forced expression of Stab2, suggesting the role of Stab2 in positive regulation of cell-cell fusion.

Stab2-deficient mice exhibited a decrease in skeletal muscle weight relative body weight, which was due to the reduction of myofiber size rather than changes in the number of myofibers. Myonuclei number per myofiber was decreased in Stab2-deficient muscle compared with wild-type muscle, indicating defective myoblast fusion by Stab2 deficiency. Consistent with this, the fusion of Stab2-deficient myoblasts was diminished than that of wild-type myoblasts. Ablation of Stab2 caused deterioration of muscle regeneration after muscle injury; the regenerating muscles of mice lacking the Stab2 gene contained smaller myofibers and fewer centralized nuclei. However, proliferation and activation of satellite cells during muscle regeneration was not affected by Stab2 ablation, which implied the role of Stab2 in myoblast fusion, but not in myogenic differentiation.

Other than as a PS receptor, Stab2 has another function as a scavenger receptor for hyaluronic acid and chondroitin sulfates. We found that a significant reduction of myoblast fusion was shown by the sole treatment of EGF-like domain repeat for PS binding and not the Link domain for hyaluronic acid and chondroitin sulfates, indicating the role of Stab2 as a PS-receptor in myoblast fusion. The blocking of external PS using anti-PS antibodies reduced myoblast fusion on wild-type myoblasts while it did not change the fusion index on Stab2-deficient myoblasts. Besides, in contrast to promotion of wild-type myoblast fusion by PS/PC liposomes, treatment with PS/PC liposomes did not raise the fusion index of Stab2 deficient myoblast. These evidences suggested that fusion-increasing activity of Stab2 was reasonably related with its function in recognition of PS exposed in myoblasts. Furthermore, excess

myonuclear accretion in C2C12/Stab2 cells was abolished by anti-PS antibodies but not by caspase inhibitors, indicating that Stab2 promotes myoblast fusion by interacting with PS exposed on live myoblasts rather than apoptotic myoblasts.

Stab2 also acts as a receptor for apoptotic cell clearance (efferocytosis). After recognizing PS exposed on apoptotic cells, Stab2 activates signaling cascades through both Gulp/Rac1 and ELMO/DOCK180/Rac1 pathways, thereby inducing actin polymerization for making the phagocytic cup (Kim et al. (2012) Mol. Cell. Biol. 32(14): 2698-2708, doi: 10.1128/MCB.06743-11). In fact, actin cytoskeleton regulator proteins, such as ELMO/Dock1, Arf-6 and N-WASP, function for myoblast fusion as well as efferocytosis, and as stated in a recent study of *C. elegans*, phagocytic pathway components including phosphatidylserine receptor (PSR-1) are also needed for regenerative axonal fusion (Neumann et al. (2015) Nature 517:219-222, doi:10.1038/nature14102). Therefore, this raised the possibility that the signal from Stab2-PS binding can recruit fusion-mediating molecules and proceed to myoblast fusion. In comparison with defects observed in Rac and Myomaker deficiencies, mild fusion phenotypes in Stab2-deficient mice may be explained by redundancy of signaling pathways. In our perspective, Stab2 and PS binding may work as a signaling cascade for “fuse-me” by recruitment of fusion mediating proteins during myofibers formation.

ACKNOWLEDGEMENTS

This study was supported by the KIST Institutional Program (Project No. 2E26320).

Keywords: Myoblast fusion, Muscle regeneration, stabilin-2, phosphatidylserine

Abbreviations: Stab2, stabilin-2; PS, phosphatidylserine

Perspective to: Seung-Yoon Park et al (2016), Stabilin-2 modulates the efficiency of myoblast fusion during myogenic differentiation and muscle regeneration, Nature Communications, 7 :10871, doi:10.1038/ncomms10871