

BMB Reports – Manuscript Submission

Manuscript Draft

Manuscript Number: BMB-20-270

Title: The Golgi complex: a hub for the secretory pathway

Article Type: Mini Review

Keywords: Golgi complex; Membrane trafficking; glycosylation; signaling transduction; organelle homeostasis

Corresponding Author: Seung-Yeol Park

Authors: Kunyou Park¹, Sung-Eun Ju¹, Nari Kim¹, Seung-Yeol Park^{1,*}

Institution: ¹Department of Life Sciences, Pohang University of Science and Technology, Pohang, Gyeongbuk, 37673 Republic of Korea,

The Golgi complex: a hub for the secretory pathway

Kunyou Park¹, Sungeun Ju¹, Nari Kim¹, Seung-Yeol Park^{1,*}

¹*Department of Life Sciences, Pohang University of Science and Technology, Pohang,
Gyeongbuk, 37673 Republic of Korea*

*Address correspondence to:

Seung-Yeol Park

Tel: 82-54-279-2325

Email: seungpark@postech.ac.kr

Abstract

The Golgi complex plays a central role in protein secretion by regulating cargo sorting and trafficking. As these processes are of functional importance to cell polarity, motility, growth, and division, there is considerable interest in achieving a comprehensive understanding of Golgi complex biology. However, the unique stack structure of this organelle has been a major hurdle to our understanding of how proteins are secreted through the Golgi apparatus. Herein, we summarize available relevant research to gain an understanding of protein secretion via the Golgi complex. This includes the molecular mechanisms of intra-Golgi trafficking and cargo export in the trans-Golgi network. Moreover, we review recent insights on signaling pathways regulated by the Golgi complex and their physiological significance.

Introduction

The intracellular transport system shuttles cargo (e.g., proteins and lipids) between organelles via transport carriers, including vesicles and tubules. Fundamentally, cargo is sorted into transport carriers generated from the membrane of donor organelles (1, 2). The cargo-containing carriers are then translocated to acceptor organelles along the microtubules (3). Upon synthesis, cargoes, which pass quality control in the endoplasmic reticulum (ER), are transported to the Golgi complex in COPII vesicles (4, 5). Thereafter, protein cargoes undergo post-translational modifications as they pass through the Golgi complex. Functional alterations in the Golgi complex have been described in different conditions, including various cancers, immunodeficiencies, and neurodegenerative diseases (6). Further, the question of how this organelle is regulated in pathological conditions has attracted considerable attention (7, 8). However, investigating the Golgi complex has been challenging due to technical limitations arising from the functional complexity and dynamic morphological changes of this organelle. In this short review, we summarize advances in the field of membrane biology from the past years and introduce the current issues in the field to obtain a better understanding of the role of the Golgi complex in the secretory pathway of cells.

Coordinated regulation of bidirectional transport within the Golgi complex

Secretory proteins undergo various post-translational modifications, including glycosylation, during their passage through the Golgi complex. How proteins are transported through this organelle has remained elusive for many years. Early studies identified the coat protein I (COPI) complex, which acts as a major coat component of vesicles formed in the Golgi (9). Extensive research pursued understanding the role of COPI vesicles in the Golgi complex. Considering the Golgi stacks as static, it was initially thought that COPI-coated vesicles are responsible for the anterograde and retrograde transport of cargoes within the organelle (10). However, this model could not explain the sorting of large cargoes such as procollagen into COPI vesicles with a small diameter of approximately 50 nm (11, 12). Subsequent studies revealed that procollagen is transported through the Golgi complex without leaving the cisternae (13, 14). Moreover, two independent research groups observed the change of cis-stack to trans-stack via live imaging in yeast (15, 16). These findings provided strong evidence for a possible mechanism through which stacks containing secretory cargoes move in the anterograde direction, referred to as the model of cisternal maturation (Fig. 1a). Given that COPI vesicles contain Golgi resident proteins, including mannosidase II and giantin (17), it was suggested that secretory cargo is transported through the Golgi complex in an anterograde direction via cisternal maturation, while the rearrangement of Golgi resident proteins is regulated by COPI vesicle-mediated retrograde transport.

COPI is a large protein complex consisting of seven subunits (α , β , β' , γ , δ , ϵ , and ξ) (18, 19). It is structurally similar to clathrin and the AP2 complex, having an inner coat (β , γ , δ , ξ), which corresponds to the AP2 adaptor, and an outer coat (α , β' , ϵ) corresponding to the clathrin heavy chain (20). Coat proteins carry out two major roles, namely cargo sorting and vesicle formation. With regard to cargo sorting, the COPI complex contains several cargo binding sites (21, 22). Structural examination revealed that KKxx and KxKxx motifs in the cytoplasmic tail of retrograde cargo are inserted into the pores of the β' -COP subunit for binding (23). Cargo binding sites were additionally identified in the α -, β' -, and γ -subunits, suggesting that cargoes bind to COPI subunits in different manners (24, 25). For instance, many Golgi resident proteins do not contain the canonical KKxx motif (26). In this case, cargoes are recognized by GOLPH3, which is a linker between Golgi-resident proteins and the COPI complex (27, 28). Indeed, various glycosyltransferases bind to GOLPH3 through a pentameric (F/L)-(L/I/V)-X-X-(R/K) motif for the COPI interaction (27, 28).

The molecular mechanisms underlying the regulation of COPI vesicle formation are unclear. It was initially suggested that small GTPases ARF1 and COPI are sufficient to form COPI vesicles based on the observation that vesicles could be reconstituted using an *in vitro* reconstitution assay wherein an intact Golgi membrane was incubated with purified COPI and recombinant ARF1 loaded with GTP γ S (29). In this scenario, the COPI complex was recruited to the Golgi membrane upon activation of ARF1 (30). The GTPase-activating protein (GAP) for ARF1, ARFGAP1, then promoted the dissociation of COPI components from vesicles via GTP hydrolysis (29). However, given that the GAP acts as a coat component of vesicle formation in multiple transport pathways (31), various groups have questioned the role of ARFGAP1 in COPI vesicle formation (32, 33). By establishing a two-stage *in vitro* reconstitution system that separates the early (budding) and late (fission) stages of COPI vesicle formation, ARFGAP1 was found to be involved in the early stage, regulating cargo sorting (34). Additionally, ARFGAP1 bound to a highly curved membrane through the ALPS motif. Further, this binding induced the membrane curvature required for vesicle formation (32, 35, 36). Taken together, these results indicated that ARFGAP1 acts as a coat component in COPI vesicle formation (36, 37). Through cryo-EM, recent studies found irregular cage structures of COPI vesicles which were reconstituted using a coatomer and ARF1 (38). Considering that vesicles coated with COPII and clathrin have a regular cage structure, COPI vesicles of different structures may be observed through the addition of missing coat components.

Although there are many advantages of the cisternal maturation model in explaining anterograde transport in the Golgi, it is still unclear why some cargoes are transported faster than others (39). As rapidly growing cells produce and secrete various molecules within a short period of time, understanding the molecular mechanisms underlying rapid secretion may be of great relevance to pathological conditions as well as normal physiology. Golgi stacks were found to be rapidly partitioned, providing different transport pathways for rapid secretion (Fig. 1b) (40). It was also suggested that certain cargoes use a special transport carrier for rapid secretion. Historically, rapidly secreted cargoes (e.g., VSV-G and proinsulin) were originally observed in COPI-coated transport carriers (10). However, this was not further investigated as the COPI complex was thought to be involved in retrograde transport. This issue was recently revisited using electron microscopy (EM) tomography, in which Z-stack images of the Golgi complex obtained via EM were applied for 3D-reconstruction. A Golgi structure in which stacks are connected by tubules was observed (41). These tubules were highly likely to be considered as vesicles in previous studies as horizontal images of the two were similar (41, 42). Subsequent studies revealed that the COPI complex promoted the formation of Golgi tubules (Fig. 1c). Mechanistically, after initial budding of the Golgi membrane induced by the COPI complex, the enzymatic activities of lysophosphatidic acid acyltransferase- γ (LPAAT γ) and cytosolic phospholipase A2 α (cPLA2 α)

determined the fate of these buds in becoming vesicles or tubules (43). In particular, cPLA2 α induced the elongation of COPI-coated buds to form tubules by providing lysophosphatidic acids with an inverted cone shape, which inhibit COPI vesicle fission (43). In this regard, rapid protein secretion was abolished by disrupting cPLA2 α , suggesting that the Golgi tubule is important for the rapid secretory pathway (43). Taken together, secretory proteins are transported through the Golgi complex by cisternal maturation (slow) and Golgi tubules (rapid) formed by the COPI complex, whereas rearrangement of Golgi resident proteins is regulated by retrograde transport via COPI vesicles.

How the same coat protein acts in the opposite direction during transport was recently revealed, as anterograde cargoes were found to bind to the COPI complex through a non-canonical basic sorting motif, distinct from retrograde cargo binding to the COPI subunit via the KKxx motif (44). This process is involved in the role of Rho GTPase CDC42 within the Golgi complex. Upon activation, CDC42 binds to the γ -subunit of the COPI complex through the KKxx motif at its C-terminus region, which causes the competitive inhibition of retrograde cargo binding to the COPI complex (25, 44). CDC42 also promotes anterograde transport through its ability to exert membrane curvature, which induces the formation of Golgi tubules (44). In this case, however, CDC42 did not affect anterograde cargo sorting into the tubules. These findings raised questions regarding how the role of CDC42 in Golgi transport is regulated. One possible mechanism is that GTPases are modulated by their respective GEF/GAP. Of note, various GEFs and GAPs of the Rho GTPase have been found in the Golgi complex (45, 46). In addition to the general hypothesis presented above, upstream kinases can target GEF/GAP to regulate GTPases as studies on Vav proteins, which are prototypic GEFs for Rho GTPases, have shown (47). The SRC kinase is a likely upstream regulator of CDC42 based on the observation that protein secretion through the Golgi complex is inhibited by SRC disruption (48). Taken together, these considerations suggest the high possibility that SRC targets the GEF/GAP for Golgi CDC42 to modulate the latter's role in transport. Studies on the circuit of SRC-GEF/GAP-Rho GTPase in Golgi transport are necessary in order to achieve a complete understanding of bidirectional transport within the organelle. Moreover, investigating the involvement of other Rho GTPases, such as Rac1 and RhoA, in the secretory pathway is another intriguing topic the elucidation of which will contribute to a more complete understanding of secretion.

Cargo sorting and transport in the trans-Golgi network

When secretory proteins arrive at the trans-Golgi network (TGN), they must be properly sorted before being delivered to their final destinations (49). TGN export was previously considered a constitutive process (50). However, cumulative studies have revealed that this pathway is regulated (Fig. 2). In fact, polarized distribution of apical and basolateral proteins provides strong evidence for the regulation of transport in the TGN. Impaired sorting causes mislocalization of basolateral cargoes in the apical region (51). Basolateral proteins (e.g., Furin and LDLR) bind to the μ subunit of the AP complex through a tyrosine-based motif (YXX Φ) and a di-leucine motif ([DE]XXXL[L]), promoting cargo export in the TGN (52). Impaired sorting of basolateral proteins by disruption of AP-4 μ subunit leads to mislocalization to the apical surface (49, 53). Cargo receptors also recognize ligands other than the peptide-based sorting motif. When apical-targeted proteins reach the TGN, the sugars in cargoes are recognized by lectins, which promotes rearrangement of apical cargoes on lipid rafts for the sorting process (54-56). The mannose-6-phosphate receptor (M6PR) also recognizes sugars on lysosomal enzymes (57). Upon cargo binding, the receptor binds to GGA and AP-1 to sort cargoes into clathrin-coated vesicles. Subsequently, low pH in the endosomal compartments induces the dissociation of cargoes from M6PR, which are then retrieved to the TGN (57). Notably, many proteins (e.g., insulin-

like growth factor II (IGF II), retinoic acid, soluble enzymes, as well as lysosomal membrane proteins) are transported independently of M6PR (52). In this regard, an alternative mechanism regulated by sortilin and LIMP-2 was identified. Sortilin contains a VPS10 domain on the lumen side. The cleavage of the N-terminal propeptide on this VPS10 domain regulates the sorting of lysosome targeting proteins such as cathepsin D and cathepsin H. Further, the cytoplasmic tail of sortilin binds to the GGA and AP complex for TGN export (58, 59). LIMP-2 is a key regulator of β -glucocerebrosidase (β GC) transport in the TGN (60). A heavily glycosylated coiled-coil domain of LIMP-2 binds to β GC. Binding is disrupted at the acidic pH once cargo is delivered to the endosomal compartment. How soluble proteins such as cartilage oligomeric protein (COMP) and lysozyme C (LyzC) are sorted after exiting the TGN was recently elucidated. That is, ADF/Cofilin1 was shown to increase calcium concentration in the lumen of the TGN through a calcium pump, SPCA1 (61). Higher Ca^{2+} promoted the oligomerization of Cab45, inducing the binding to soluble cargoes for TGN export (62). Although extensive studies have uncovered regulatory factors of TGN export, the coat protein that regulates various transport pathways occurring in the TGN remains unclear.

Signaling pathways regulated by the Golgi complex for organelle homeostasis

The Golgi complex is involved in the activation of signaling transduction by providing a platform for signaling molecules (Fig. 3). For example, the signaling of N-Ras in the Golgi complex increases T cell responses to antigen stimulation (63, 64). Ras signaling initiated from the Golgi complex triggers sustained T cell activity compared to that from the plasma membrane (65). These observations lead to the question of how signaling molecules are specifically recruited to the Golgi complex, but not to other organelles. The membrane association of small GTPases was described as regulated by lipid modifications, including farnesylation, prenylation, and palmitoylation (66). Further, the recruitment of Ras to the Golgi complex was determined by a balance between palmitoylation and depalmitoylation (67, 68). Given that many upstream regulators, such as GEF and GAP, exist in the Golgi complex (45, 46), it is possible that these are initially recruited to the apparatus to activate the Rho GTPase on its membrane. Notably, however, how GEF and GAP are recruited to specific organelles and what signaling activates Rho GTPase in TGN export are both poorly understood concepts.

Adenylyl cyclase, which converts cellular ATP to cyclic AMP (cAMP), regulates PKA signaling upon activation of G-protein coupled receptors (GPCRs) (69, 70). PKA promotes the recruitment of ARF1 to the Golgi complex, resulting in membrane trafficking (71). Since GPCR signaling is mostly regulated at the plasma membrane, it is surprising that the Golgi complex is involved. In this regard, recent studies have uncovered a new role of the KDEL receptor (KDELR), a GPCR, in signaling transduction (72, 73). KDELR was originally known as a cargo receptor for COPI transport. Under ER stress conditions, ER resident proteins often leak from the ER and are captured by the KDELR located at the Golgi complex, which binds to the "KDEL" tetrapeptide sequence of ER resident proteins. This promotes the retrieval of proteins back to the ER through COPI vesicles (74, 75). Studies investigating its physiological role revealed that different signaling pathways activated by KDELR regulate distinct transport pathways, including Gq/SRC for cargo export at the TGN as well as Gs/PKA for COPI transport (72, 73). This raises the key question regarding how various signaling pathways are regulated by this single receptor. Among the three types of KDELR (KDELR1/2/3) in mammalian cells, only KDELR1-mediated signaling has been studied. We suspect that the multitude of different G-proteins coupled to KDELRs are likely to activate different downstream effector proteins. A proteomic approach using engineered ascorbic acid peroxidase (APEX) was recently utilized to determine the spatiotemporal protein networks of GPCRs in living cells (76, 77). This could be an appropriate

technical approach for identifying new pathways regulated by KDEL_R in the Golgi complex.

Intracellular compartments maintain organelle homeostasis in response to environmental stress through a number of mechanisms. Over the past decades, the endoplasmic reticulum (ER) has been extensively studied with regard to the maintenance of homeostasis under stress. Protein synthesis is considerably enhanced under various physiological and pathological conditions. Such increased rates of protein synthesis overwhelm the protein folding capacity of the ER, leading to the accumulation of misfolded protein (78). Multiple stress response mechanisms are activated to maintain organelle homeostasis and are cumulatively known as the unfolded protein response (UPR). As proteins are secreted through the Golgi, how it maintains homeostasis in the context of increased transport flux is another topic of considerable interest. However, a limited number of studies have the Golgi complex stress response. In this regard, proteins such as TFE3, proteoglycans, CREB3, and HSP47 have been identified as key factors in the maintenance Golgi homeostasis (79, 80). However, the mechanisms through which these factors regulate the Golgi stress response remain unclear. Further, since the Golgi is involved in various signaling pathways, it is likely that KDEL_R-associated signaling pathways promote transport through the organelle, reducing protein amounts within. Given the complex dynamics of the Golgi complex in diverse conditions, further research is needed to elucidate aspects such as the Golgi stress response and related signaling pathways.

CONCLUSION

The Golgi complex plays an important role in various cellular processes. Extensive studies have been conducted in order to understand the organelle from a membrane trafficking perspective. However, Golgi regulation under different conditions remains unclear. In addition to studies on membrane trafficking and related signaling, there are various fundamental questions that need to be addressed regarding the Golgi complex, such as the identification of signaling cues, understanding how organelle homeostasis is maintained, and elucidating the significance of glycosylation in the context of membrane trafficking. Moreover, a new form of intracellular communication regulated by direct organelle contact was recently discovered. This raises the question of whether the Golgi complex also communicates with other organelles through membrane contact and, if so, what the physiological significance of Golgi contact is.

ACKNOWLEDGEMENTS

This study was supported by grants from the National Research Foundation (NRF-2020R1C1C100882311, NRF-2017R1A5A101536614), the Korea Health Industry Development Institute (KHIDI-HR20C0025), POSCO Science Fellowship, and the BK21 Plus and BK21 FOUR Research Fellowship.

CONFLICTS OF INTEREST

The authors have no conflicting interests relevant to this study to disclose.

REFERENCES

1. Sato K and Nakano A (2007) Mechanisms of COPII vesicle formation and protein sorting. *FEBS Letters* 581, 2076-2082
2. Hsu VW, Lee SY and Yang JS (2009) The evolving understanding of COPI vesicle formation. *Nat Rev Mol Cell Biol* 10, 360-364
3. Fourriere L, Jimenez AJ, Perez F and Boncompain G (2020) The role of microtubules in secretory protein transport. *Journal of Cell Science* 133, jcs237016
4. Barlowe C, Orci L, Yeung T et al (1994) COPII: a membrane coat formed by Sec proteins that drive vesicle budding from the endoplasmic reticulum. *Cell* 77, 895-907
5. Zanetti G, Pahuja KB, Studer S, Shim S and Schekman R (2011) COPII and the regulation of protein sorting in mammals. *Nat Cell Biol* 14, 20-28
6. Makhoul C, Gosavi P and Gleeson PA (2019) Golgi Dynamics: The Morphology of the Mammalian Golgi Apparatus in Health and Disease. *Front Cell Dev Biol* 7, 112
7. Lee TH and Linstedt AD (1999) Osmotically induced cell volume changes alter anterograde and retrograde transport, Golgi structure, and COPI dissociation. *Mol Biol Cell* 10, 1445-1462
8. Kellokumpu S, Sormunen R and Kellokumpu I (2002) Abnormal glycosylation and altered Golgi structure in colorectal cancer: dependence on intra-Golgi pH. *FEBS Lett* 516, 217-224
9. Rothman JE and Wieland FT (1996) Protein Sorting by Transport Vesicles. *Science* 272, 227
10. Orci L, Stames M, Ravazzola M et al (1997) Bidirectional Transport by Distinct Populations of COPI-Coated Vesicles. *Cell* 90, 335-349
11. Becker B and Melkonian M (1996) The secretory pathway of protists: spatial and functional organization and evolution. *Microbiological Reviews* 60, 697
12. Bonfanti L, Mironov AA, Jr., Martínez-Menárguez JA et al (1998) Procollagen Traverses the Golgi Stack without Leaving the Lumen of Cisternae: Evidence for Cisternal Maturation. *Cell* 95, 993-1003
13. Mironov AA, Beznoussenko GV, Nicoziani P et al (2001) Small cargo proteins and large aggregates can traverse the Golgi by a common mechanism without leaving the lumen of cisternae. *Journal of Cell Biology* 155, 1225-1238
14. Donohoe BS, Kang B-H and Staehelin LA (2007) Identification and characterization of COPIa- and COPIb-type vesicle classes associated with plant and algal Golgi. *Proceedings of the National Academy of Sciences* 104, 163
15. Losev E, Reinke CA, Jellen J, Strongin DE, Bevis BJ and Glick BS (2006) Golgi maturation visualized in living yeast. *Nature* 441, 1002-1006
16. Matsuura-Tokita K, Takeuchi M, Ichihara A, Mikuriya K and Nakano A (2006) Live imaging of yeast Golgi cisternal maturation. *Nature* 441, 1007-1010
17. Martínez-Menárguez JA, Prekeris R, Oorschot VMJ et al (2001) Peri-Golgi vesicles contain retrograde but not anterograde proteins consistent with the cisternal progression model of intra-Golgi transport. *Journal of Cell Biology* 155, 1213-1224
18. Lowe M and Kreis TE (1998) Regulation of membrane traffic in animal cells by COPI. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 1404, 53-66
19. Nickel W, Brügger B and Wieland FT (2002) Vesicular transport: the core machinery of COPI recruitment and budding. *Journal of Cell Science* 115, 3235
20. Hoffman GR, Rahl PB, Collins RN and Cerione RA (2003) Conserved Structural Motifs in Intracellular Trafficking Pathways: Structure of the γ COP Appendage Domain. *Molecular Cell* 12, 615-625
21. Jackson MR, Nilsson T and Peterson PA (1990) Identification of a consensus motif for retention of transmembrane proteins in the endoplasmic reticulum. *The EMBO Journal* 9, 3153-3162
22. Cosson P and Letourneur F (1994) Coatamer interaction with di-lysine endoplasmic reticulum retention motifs. *Science* 263, 1629
23. Jackson Lauren P, Lewis M, Kent Helen M et al (2012) Molecular Basis for Recognition of Dilysine Trafficking Motifs by COPI. *Developmental Cell* 23, 1255-1262
24. Ma W and Goldberg J (2013) Rules for the recognition of dilysine retrieval motifs by coatamer. *The EMBO Journal* 32, 926-937
25. Wu WJ, Erickson JW, Lin R and Cerione RA (2000) The γ -subunit of the coatamer complex binds Cdc42 to mediate transformation. *Nature* 405, 800-804
26. Eckert ESP, Reckmann I, Hellwig A et al (2014) Golgi Phosphoprotein 3 Triggers Signal-mediated Incorporation of Glycosyltransferases into Coatamer-coated (COPI) Vesicles. *Journal of Biological Chemistry*
27. Tu L, Tai WCS, Chen L and Banfield DK (2008) Signal-Mediated Dynamic Retention of

- Glycosyltransferases in the Golgi. *Science* 321, 404
28. Dippold HC, Ng MM, Farber-Katz SE et al (2009) GOLPH3 Bridges Phosphatidylinositol-4-Phosphate and Actomyosin to Stretch and Shape the Golgi to Promote Budding. *Cell* 139, 337-351
 29. Tanigawa G, Orci L, Amherdt M, Ravazzola M, Helms JB and Rothman JE (1993) Hydrolysis of bound GTP by ARF protein triggers uncoating of Golgi-derived COP-coated vesicles. *Journal of Cell Biology* 123, 1365-1371
 30. Donaldson JG, Cassel D, Kahn RA and Klausner RD (1992) ADP-ribosylation factor, a small GTP-binding protein, is required for binding of the coatomer protein beta-COP to Golgi membranes. *Proceedings of the National Academy of Sciences* 89, 6408
 31. Kung LF, Pagant S, Futai E et al (2012) Sec24p and Sec16p cooperate to regulate the GTP cycle of the COPII coat. *The EMBO Journal* 31, 1014-1027
 32. Bigay J, Gounon P, Robineau S and Antonny B (2003) Lipid packing sensed by ArfGAP1 couples COPI coat disassembly to membrane bilayer curvature. *Nature* 426, 563-566
 33. Reinhard C, Schweikert M, Wieland FT and Nickel W (2003) Functional reconstitution of COPI coat assembly and disassembly using chemically defined components. *Proceedings of the National Academy of Sciences* 100, 8253
 34. Lanoix J, Ouwendijk J, Stark A et al (2001) Sorting of Golgi resident proteins into different subpopulations of COPI vesicles : a role for ArfGAP1. *Journal of Cell Biology* 155, 1199-1212
 35. Bigay J, Casella J-F, Drin G, Mesmin B and Antonny B (2005) ArfGAP1 responds to membrane curvature through the folding of a lipid packing sensor motif. *The EMBO Journal* 24, 2244-2253
 36. Park S-Y, Yang J-S, Li Z et al (2019) The late stage of COPI vesicle fission requires shorter forms of phosphatidic acid and diacylglycerol. *Nature Communications* 10, 3409
 37. Yang J-S, Lee SY, Gao M et al (2002) ARFGAP1 promotes the formation of COPI vesicles, suggesting function as a component of the coat. *Journal of Cell Biology* 159, 69-78
 38. Dodonova SO, Diestelkoetter-Bachert P, von Appen A et al (2015) A structure of the COPI coat and the role of coat proteins in membrane vesicle assembly. *Science* 349, 195
 39. Glick BS and Nakano A (2009) Membrane Traffic Within the Golgi Apparatus. *Annual Review of Cell and Developmental Biology* 25, 113-132
 40. Patterson GH, Hirschberg K, Polishchuk RS, Gerlich D, Phair RD and Lippincott-Schwartz J (2008) Transport through the Golgi Apparatus by Rapid Partitioning within a Two-Phase Membrane System. *Cell* 133, 1055-1067
 41. Trucco A, Polishchuk RS, Martella O et al (2004) Secretory traffic triggers the formation of tubular continuities across Golgi sub-compartments. *Nature Cell Biology* 6, 1071-1081
 42. Pietro ES, Capestrano M, Polishchuk EV et al (2009) Group IV Phospholipase A2 α Controls the Formation of Inter-Cisternal Continuities Involved in Intra-Golgi Transport. *PLOS Biology* 7, e1000194
 43. Yang J-S, Valente C, Polishchuk RS et al (2011) COPI acts in both vesicular and tubular transport. *Nature Cell Biology* 13, 996-1003
 44. Park S-Y, Yang J-S, Schmider AB, Soberman RJ and Hsu VW (2015) Coordinated regulation of bidirectional COPI transport at the Golgi by CDC42. *Nature* 521, 529
 45. Farhan H and Hsu VW (2016) Cdc42 and Cellular Polarity: Emerging Roles at the Golgi. *Trends in Cell Biology* 26, 241-248
 46. Baschieri F, Confalonieri S, Bertalot G et al (2014) Spatial control of Cdc42 signalling by a GM130-RasGRF complex regulates polarity and tumorigenesis. *Nature Communications* 5, 4839
 47. Bustelo XR (2001) Vav proteins, adaptors and cell signaling. *Oncogene* 20, 6372-6381
 48. Pulvirenti T, Giannotta M, Capestrano M et al (2008) A traffic-activated Golgi-based signalling circuit coordinates the secretory pathway. *Nature Cell Biology* 10, 912-922
 49. Guo Y, Sirkis DW and Schekman R (2014) Protein Sorting at the trans-Golgi Network. *Annual Review of Cell and Developmental Biology* 30, 169-206
 50. Polishchuk EV, Di Pentima A, Luini A and Polishchuk RS (2003) Mechanism of Constitutive Export from the Golgi: Bulk Flow via the Formation, Protrusion, and En Bloc Cleavage of large trans-Golgi Network Tubular Domains. *Molecular Biology of the Cell* 14, 4470-4485
 51. Tanos B and Rodriguez-Boulan E (2008) The epithelial polarity program: machineries involved and their hijacking by cancer. *Oncogene* 27, 6939-6957
 52. Dittmer F, Ulbrich EJ, Hafner A et al (1999) Alternative mechanisms for trafficking of lysosomal enzymes in mannose 6-phosphate receptor-deficient mice are cell type-specific. *Journal of Cell Science* 112, 1591
 53. Simmen T, Höning S, Icking A, Tikkanen R and Hunziker W (2002) AP-4 binds basolateral signals and participates in basolateral sorting in epithelial MDCK cells. *Nature Cell Biology* 4, 154-159

54. Brewer CF, Miceli MC and Baum LG (2002) Clusters, bundles, arrays and lattices: novel mechanisms for lectin–saccharide-mediated cellular interactions. *Current Opinion in Structural Biology* 12, 616-623
55. Weisz OA and Rodriguez-Boulan E (2009) Apical trafficking in epithelial cells: signals, clusters and motors. *Journal of Cell Science* 122, 4253
56. Scheiffele P, Peränen J and Simons K (1995) N-glycans as apical sorting signals in epithelial cells. *Nature* 378, 96-98
57. Kornfeld S and Mellman I (1989) The Biogenesis of Lysosomes. *Annual Review of Cell Biology* 5, 483-525
58. Lefrancois S, Zeng J, Hassan AJ, Canuel M and Morales CR (2003) The lysosomal trafficking of sphingolipid activator proteins (SAPs) is mediated by sortilin. *The EMBO Journal* 22, 6430-6437
59. Ni X and Morales CR (2006) The Lysosomal Trafficking of Acid Sphingomyelinase is Mediated by Sortilin and Mannose 6-phosphate Receptor. *Traffic* 7, 889-902
60. Reczek D, Schwake M, Schröder J et al (2007) LIMP-2 Is a Receptor for Lysosomal Mannose-6-Phosphate-Independent Targeting of β -Glucocerebrosidase. *Cell* 131, 770-783
61. von Blume J, Alleaume A-M, Cantero-Recasens G et al (2011) ADF/Cofilin Regulates Secretory Cargo Sorting at the TGN via the Ca²⁺ ATPase SPCA1. *Developmental Cell* 20, 652-662
62. von Blume J, Alleaume A-M, Kienzle C, Carreras-Sureda A, Valverde M and Malhotra V (2012) Cab45 is required for Ca²⁺-dependent secretory cargo sorting at the trans-Golgi network. *Journal of Cell Biology* 199, 1057-1066
63. Bivona TG, Quatela S and Philips MR (2006) Analysis of Ras activation in living cells with GFP-RBD. *Methods Enzymol* 407, 128-143
64. Ibiza S, Perez-Rodriguez A, Ortega A et al (2008) Endothelial nitric oxide synthase regulates N-Ras activation on the Golgi complex of antigen-stimulated T cells. *Proc Natl Acad Sci U S A* 105, 10507-10512
65. Yasuda T and Kurosaki T (2008) Regulation of lymphocyte fate by Ras/ERK signals. *Cell Cycle* 7, 3634-3640
66. Ahearn IM, Haigis K, Bar-Sagi D and Philips MR (2011) Regulating the regulator: post-translational modification of RAS. *Nat Rev Mol Cell Biol* 13, 39-51
67. Goodwin JS, Drake KR, Rogers C et al (2005) Depalmitoylated Ras traffics to and from the Golgi complex via a nonvesicular pathway. *J Cell Biol* 170, 261-272
68. Rocks O, Peyker A, Kahms M et al (2005) An acylation cycle regulates localization and activity of palmitoylated Ras isoforms. *Science* 307, 1746-1752
69. Daaka Y, Luttrell LM and Lefkowitz RJ (1997) Switching of the coupling of the β 2-adrenergic receptor to different G proteins by protein kinase A. *Nature* 390, 88-91
70. Clark RB, Knoll BJ and Barber R (1999) Partial agonists and G protein-coupled receptor desensitization. *Trends in Pharmacological Sciences* 20, 279-286
71. Muniz M, Martin ME, Hidalgo J and Velasco A (1997) Protein kinase A activity is required for the budding of constitutive transport vesicles from the trans-Golgi network. *Proc Natl Acad Sci U S A* 94, 14461-14466
72. Cancino J, Capalbo A, Di Campli A et al (2014) Control Systems of Membrane Transport at the Interface between the Endoplasmic Reticulum and the Golgi. *Developmental Cell* 30, 280-294
73. Giannotta M, Ruggiero C, Grossi M et al (2012) The KDEL receptor couples to G α q&11 to activate Src kinases and regulate transport through the Golgi. *The EMBO Journal* 31, 2869
74. Semenza JC, Hardwick KG, Dean N and Pelham HRB (1990) ERD2, a yeast gene required for the receptor-mediated retrieval of luminal ER proteins from the secretory pathway. *Cell* 61, 1349-1357
75. Lewis MJ and Pelham HRB (1992) Ligand-induced redistribution of a human KDEL receptor from the Golgi complex to the endoplasmic reticulum. *Cell* 68, 353-364
76. Paek J, Kalocsay M, Staus DP et al (2017) Multidimensional Tracking of GPCR Signaling via Peroxidase-Catalyzed Proximity Labeling. *Cell* 169, 338-349.e311
77. Lobingier BT, Hüttenhain R, Eichel K et al (2017) An Approach to Spatiotemporally Resolve Protein Interaction Networks in Living Cells. *Cell* 169, 350-360.e312
78. Bernales S, Papa FR and Walter P (2006) Intracellular Signaling by the Unfolded Protein Response. *Annual Review of Cell and Developmental Biology* 22, 487-508
79. Sun Z and Brodsky JL (2019) Protein quality control in the secretory pathway. *Journal of Cell Biology* 218, 3171-3187
80. Sasaki K and Yoshida H (2019) Golgi stress response and organelle zones. *FEBS Letters* 593, 2330-2340

FIGURE LEGNEDS**Figure1. Three models of protein transport within the Golgi complex.**

(a) Cisternal maturation model. Secretory cargo is transported in an anterograde direction along with cisternae maturation from cis-face to trans-face of the Golgi complex. Golgi resident proteins are transported in retrograde through COPI vesicles. (b) Rapid partitioning model. The Golgi stacks are distributed into two sub-domains. One is glycerophospholipids containing processing domain (white), while the other domain contains glycosphingolipids (black). Transmembrane secretory protein (green) are concentrated at export domain, while Golgi-resident proteins (red) are excluded from export domain and concentrated in processing domain. (c) Golgi tubule-mediated transport model. Each cisternae is connected by Golgi tubule which is regulated by the COPI complex. Golgi tubules are involve in the rapid bidirectional transport within the Golgi complex.

Figure 2. Cargo sorting and export regulated in the trans-Golgi network.

Transport of cargoes to various destinations is regulated in the TGN. Many key factors have been identified. TGN export to endo-lysosome is regulated through Clathrin dependent and independent machineries. Mannose-6-phosphasphate receptor (M6PR) and Sortilin act as a cargo receptor in this pathway. In polarized cells, transport of apical and basolateral cargoes is regulated by multiple mechanisms that involve not only proteins, but also sugars.

Figure 3. Signaling transduction instigated by the Golgi complex.

(Left) Palmitoylation of Ras obtained at the Golgi complex is required for membrane recruitment and activation of signaling pathways. Signaling pathway of N-Ras at the Golgi complex shows a sustained activity as compared to N-Ras signaling instigated from the plasma membrane. (Right) Retrograde COPI transport is regulated by KDELR-mediated PKA signaling pathway. Cargo export in TGN is regulated by SRC signaling pathway which is activated by KDELR.

Fig 1




Fig 2




Fig 3

