

BMB Reports – Manuscript Submission

Manuscript Draft

Manuscript Number: BMB-16-141

Title: Extra-gonadal sites of estrogen biosynthesis and function

Article Type: Mini Review

Keywords: Extra-gonadal; Estrogen; Estrogen receptor; immune system; cancer development

Corresponding Author: CheMyong Jay Ko

Authors: Radwa Barakat¹, Oliver Oakley¹, Heehyen Kim¹, Jooyoung Jin¹, CheMyong Jay Ko^{1,*}

Institution: ¹Department of Comparative Biosciences, University of Illinois at Urbana-Champaign, Illinois, United States,

²Department of Toxicology, Faculty of Veterinary Medicine, Benha University, Egypt,

³Department of Biology, College of Arts and Sciences, Eastern Kentucky University, Kentucky, United States,

⁴Department of Agricultural Biotechnology, Seoul National University, Seoul, Korea,

Manuscript Type: Mini Review

Title: Extra-gonadal sites of estrogen biosynthesis and function

Radwa Barakat^{1,2}, Oliver Oakley³, Heehyen Kim⁴, Jooyoung Jin⁴ and CheMyong Jay Ko¹

¹Department of Comparative Biosciences, University of Illinois at Urbana-Champaign, Illinois, United States; ²Department of Toxicology, Faculty of Veterinary Medicine, Benha University, Egypt. ³Department of Biology, College of Arts and Sciences, Eastern Kentucky University, Kentucky, United States; ⁴Department of Agricultural Biotechnology, Seoul National University, Seoul, Korea

Running Title: non-gonadal estrogen

Keywords: Extra-gonadal, Estrogen, Estrogen receptor

Corresponding Author's Information:

CheMyong Jay Ko, PhD

Tel: 217-333-9362

Email: jayko@illinois.edu

Department of Comparative Biosciences

College of Veterinary Medicine

University of Illinois at Urbana-Champaign

3806 VMBSB, MC-002

2001 South Lincoln Avenue

Urbana, Illinois 61802, USA

Abstract

Estrogens are the key hormones regulating the development and function of reproductive organs in all vertebrates. Recent evidence indicates that estrogens play important roles in the immune system, cancer development, and other critical biological processes related to human well-being [1-4]. Obviously, the gonads (ovary and testis) are the primary sites of estrogen synthesis, but estrogens synthesized in extra-gonadal sites play an equally important role in controlling biological activities [5-7]. Understanding non-gonadal sites of estrogen synthesis and function is crucial and will lead to therapeutic interventions targeting estrogen signaling in disease prevention and treatment. Developing a rationale targeting strategy remains challenging because knowledge of extra-gonadal biosynthesis of estrogens, and the mechanism by which estrogen activity is exerted, is very limited. In this review, we will summarize recent discoveries of extra-gonadal sites of estrogen biosynthesis and their local functions and discuss the significance of the most recent novel discovery of intestinal estrogen biosynthesis.

Introduction. Estrogens are a class of steroid hormones that regulate the development and function of male and female reproductive organs. In the ovary, estrogen synthesis begins in theca cells with androgen synthesis and ends with conversion of androgens to estrogens in granulosa cells by the enzyme aromatase. In the male gonad, estrogens are synthesized in the Leydig cells, Sertoli cells, and mature spermatocytes [8]. Like other steroid hormones, estrogens enter passively into the cells and bind to the estrogen receptors, which then regulate the transcription of downstream estrogen-responsive genes. Among the number of different forms of estrogens, 17β -estradiol (estradiol) is the most common and potent form of estrogen in mammals. Estradiol is also produced in a number of extra-gonadal organs, including the adrenal glands, brain, adipose tissue, skin, pancreas [5-7], and other sites yet to be identified. The discoveries of extra-gonadal sites of estradiol synthesis greatly expands our knowledge of the novel roles of estrogens beyond the reproductive system.

Extra-gonadal sites of estrogen synthesis and its local roles. The first discovery of extra-gonadal estrogen synthesis was made in 1974 by Hemsell and his colleagues when they made an unexpected observation that androgens were converted to estrogens in adipose tissue [9]. Since then, a number of other extra-gonadal sites of estrogen synthesis have been discovered. Adipose tissues are considered to be the major source of circulating estrogen after the gonads in both men and women, and the contribution made by the adipose tissues to the total circulating estrogens increases with advancing age [9]. The chemical structure and biological activity of the estrogens synthesized in the extra-gonadal sites are not different from those that are produced by the gonads. However, there are unique features that make the extra-gonadal estrogen synthesis differ from the gonadal synthesis. A major difference is in the biochemical pathway of estrogen synthesis. The tissues and cells of the extra-gonadal sites of estrogen synthesis are

unable to synthesize C19 steroids, the precursors of estrogen synthesis, but are able to convert C19 steroids to estrogens, a critical and rate-limiting step mediated by Cyp19 aromatase. Hence, extra-gonadal estrogen synthesis is dependent on an external source of C19 precursors [7] and the level of aromatase expression. Because C19 steroids can be supplied to a local tissue via circulation and are converted to estrogens in any tissue where aromatase is expressed, the presence of aromatase expression in a local tissue confirms extra-gonadal estrogen synthesis. **Table 1** lists the peripheral tissues that express aromatase and are therefore able to convert C19 precursors to estrogens. These extra-gonadally synthesized estrogens are thought to act and be metabolized locally, which limits their systemic effects [10]. Another unique feature of extra-gonadal estrogen synthesis is that while the total amount of estrogen synthesized in each tissue may be small, the local tissue concentrations of estrogens could be high enough to exert biological impact locally. The functional roles of estrogens are mediated mostly by estrogen receptors that are nuclear receptor transcription factors. Therefore, a tissue that expresses one or more estrogen receptors is considered to be a target of estrogenic regulation. **Table 2** lists key organs and tissues that express estrogen receptors.

1. Extra-gonadal sites of estrogen synthesis

Sites	Evidence of 17 β -estradiol synthesis			References
	Cyp19 mRNA	Cyp19 protein	17 β -estradiol	
Brain	Astrocyte (rat, mouse, human), Hippocampus and hypothalamus (rat, mouse, monkey, human)	Astrocyte (mouse), GnRH (rat), Dentate gyrus/ pyramidal cell (rat, mouse, human, monkey), Interneurons(human), Granular cell (human, monkey), Purkinje cell (human, mouse), Ependymal and subependymal cell (human).	Astrocyte (rat, monkey).	[11-20]
Fat	Stromal cell (human), Adipocyte (human)	Stromal cell (human), Adipocyte, mesenchymal cell (human)		[17-20]
Bone	Osteoblast (human)	Osteoblast (human)	Osteoblast (human)	[21-23]
Liver	HepG2 hepatoma and hepatocellular carcinoma (human), Hepatocyte (porcine).	HepG2 hepatoma and hepatocellular carcinoma (human)	HepG2 hepatoma and hepatocellular carcinoma (human)	[24-26]
Adrenal gland	Adrenocortical cell (human, porcine, rat)	Adrenocortical cell (human)	Adrenocortical cell (rat)	[26-28]
Intestine	Parietal cell (rat)	Parietal cell (rat)	Parietal cell (rat)	[29]
Skin	Fibroblast (human). Keratinocyte (human).	Fibroblast (chicken, human), Keratinocyte (human)	Fibroblast (chicken)	[30-32]
Blood vessel	Smooth muscle cell (human, rat, bovine)	Smooth muscle cell (human, rat, bovine)		[33-35]
Spleen	T cell (mouse)	T cell (mouse)		[36]

2. Extra-gonadal sites of estrogen receptor expression

Sites	Receptor subtypes			References
	ER α	ER β	Other receptors	
Brain	Cholinergic neuron (rat), GABAergic neuron (rat), Pro-opiomelanocortin neuron (mouse).	GnRH neurons (mouse), Subiculum neuron (monkey), Ammon's horn neuron (monkey)	GPER1 (glial cell, rat), GPER1 (GABAergic neuron, rat).	[37-41]
Fat	Adipocyte (human)	Adipocyte (human)	GPER (adipocyte, mouse)	[42-44]
Bone	Osteoblast (mouse), Osteocyte (mouse).	Osteoblast (human), Osteocyte (human), Osteoclast (rat, human).		[45-50]
Liver	Hepatocyte (rat)			[29]
Blood vessel	Smooth muscle cell (human), Vascular endothelial cell (human)	Endothelial cell (human)	GPR30 (endothelial cell, rat)	[51-54]
Intestine	Epithelial cell (rat), Parietal cell (rat), Myenteric neuron (rat)	Epithelial cell (rat), Parietal cell (rat)	GPER (colonic epithelia, human)	[55, 56]
Skin	Keratinocyte (human), Mast cell (human) Sebocyte (human)	Keratinocyte (human), Mast cell (human)		[57-59]
Adrenal gland	Adrenal cortex (rat)	Adrenal cortex (rat)	GPER1 (rat)	[60]
Muscle	Satellite cell (rat)	Satellite cell (rat)		[61]
Kidney	Mesangial cells (human, mouse)	Mesangial cells (human, mouse)		[62]
Pancreas	β -cell (mouse)	β -cell (mouse)		[63, 64]

Adipose tissues. Adipose tissues, where estradiol stimulates the production of high density lipoprotein cholesterol (HDL) and triglycerides while decreasing LDL production and and fat deposition [65, 66], are the most extensively studied sites of extra-gonadal estrogen synthesis. Both male and female aromatase-

deficient (Cyp19KO) mice exhibit obesity and dyslipidemia [65, 66], proving that estradiol plays a beneficial role in the lipogenesis. However, an adverse effect of adipose tissue-driven estradiol is also indicated in the pathogenesis of breast cancer. For instance, in a breast with a tumor, adipose tissues proximal to the tumor exhibit higher aromatase activity than those distal to the tumor [67].

Bone. Aromatase expression in human bone has been demonstrated in osteoblasts, chondrocytes, and fibroblasts (**Table 1**), where they convert circulating androgens into estrogens [68]. In the bone of prepubertal children, the locally synthesized estradiol stimulates epiphyseal maturation during the growth phase [69]. However, in both males and females, the massive pubertal increase of estradiol leads to increased apoptosis of chondrocytes in the epiphyseal plate, causing chondrocyte depletion and hence, ossification and growth slow-down [70]. In adults, estradiol increases bone formation and mineralization and reduces bone resorption, thus reducing the risk of osteoporosis (68). Therefore, it is not surprising that the incidence of osteoporosis increases in postmenopausal women as their ovaries lose estradiol synthetic capacity.

Skin. Aromatase expression in the skin occurs mainly in hair follicles and sebaceous glands [71]. Glucocorticoids, cAMP analogs, growth factors, and cytokines modulate aromatase expression in these cells and therefore, local estrogen synthesis [72]. Estradiol enhances collagen synthesis, increases skin thickness, and stimulates blood flow in the skin. Therefore, in situ estrogen synthesis in the skin is vital for maintaining healthy skin [73]. Estradiol also prolongs the anagen phase of the hair cycle and therefore enhances hair growth by increasing the synthesis of essential growth factors stimulating the proliferation of hair follicle cells [74].

Liver. In the liver, estradiol regulates protein synthesis, including lipoprotein and proteins responsible for blood clotting (factors II, VII, IX, X, plasminogen) [75]. Estrogen signaling is also essential in regulating glucose homeostasis, thus improving glucose tolerance and insulin sensitivity [76]. Recent research has explored the possibility that postmenopausal women with nonalcoholic fatty liver disease and with long durations of estrogen deficiency could have a higher risk of having severe fibrosis than premenopausal women [77]. Estrogen receptor beta (ER β) is implicated in mediating the protective role that estradiol plays under pathogenic condition in the liver as it shows potent anti-proliferative and anti-inflammatory properties. As such, chronic disease is linked to elevated ER β expression in the liver [78]. ER β is also known to mediate the anti-tumor action of estrogens in intrahepatic cholangiocarcinoma [79].

Brain. High levels of estrogen receptors are expressed during brain development. During this period, sex hormones determine apoptosis, neuronal migration, neurogenesis, axonal guidance, and synaptogenesis. Estradiol induces sexual differentiation in the developing brain. Aromatase mRNA expression in the hypothalamus of males peaks before and after birth, inducing sexual differentiation of the brain [80]. In the brains of both males and females, estradiol provides a neuroprotective effect. Estradiol's prevention of neurodegeneration in brain tissues is proven in both the Cyp19KO mouse model and the aromatase

inhibitor-treated mouse [12]. Inhibition or null mutation of aromatase, a key enzyme for estradiol synthesis, results in accelerated neurodegeneration [12]. Estradiol effects in the brain also include regulating mood, pain sensitivity, motor control, and cognitive behavior [17-20]. Estradiol regulates neuronal metabolism by modulating the expression of metabolic enzymes such as GLUT (glucose-transporter), glycolytic enzyme hexokinase, pyruvate dehydrogenase (PDH), aconitase, and ATP synthase [81].

Adrenal gland. Estrogens stimulate adrenal cortex growth during development by promoting cell proliferation and enhancing steroidogenic activity by increasing StAR and SF-1 expression in the adrenal gland [34]. In the fetal adrenal gland, estradiol and ACTH form as a positive regulatory loop in which estradiol increases ACTH secretion from adrenal cortex while ACTH increase estradiol in the ovary [82].

Pancreas. Estradiol increases insulin gene expression and insulin content in β -cells [63, 83], increases β -cell proliferation during pancreatic development and recovery from injury [84], and prevents apoptosis of β -cells upon inflammatory insult [63] via $ER\alpha$ - and $ER\beta$ -mediated pathways.

Others. In the blood vessel, estradiol positively impacts vascular function by preventing the oxidation of LDL cholesterol, stimulating nitric oxide synthesis and release, and inhibiting fibroblast transition to myofibroblast, preventing cardiac fibrosis [85-87] and atherosclerosis development. In the muscle, estradiol increases muscle mass and strength, alleviating disuse-induced muscle atrophy and promoting regrowth after reloading. It also stimulates muscle repair by stimulating satellite cell proliferation [88, 89]. Estradiol replacement on ovariectomized mice shows that estradiol can reduce stiffness in muscle as well as stimulate muscle regeneration [43]. In the kidney, estradiol has a role of protecting kidney functions during progressive glomerulosclerosis in the female rat remnant kidney model [90]. In the intestine, to maintain the intestinal epithelium, estrogens are necessary. Estrogens improve epithelial barriers and reduce intestinal permeability [91], preventing chronic mucosal inflammation in animals and humans.[92]

Inflammation. Estrogens play an important role in the inflammatory response by regulating development, proliferation, migration, and apoptosis of immune cells [93]. Lymphocytes have been shown to express estrogen receptors ($ER\alpha$ and $ER\beta$), but the expression levels of both receptors vary among cell types. $CD4^+$ T-lymphocytes express $ER\alpha$ whereas B-lymphocytes express $ER\beta$ [94]. In contrast, $CD8^+$ T-lymphocytes express both receptors at low but equivalent levels [94]. Regardless of subcellular differences, estrogens appear to exert a suppressive effect on both B- and T-lymphopoiesis. In support, B-lymphocyte formation is selectively reduced with estradiol treatment [95], and ovariectomy results in increased B-lymphopoiesis [96, 97]. In addition to the inhibitory effect on lymphopoiesis, estradiol has been shown to influence T helper (Th) responses; inhibit the production of Th1 cytokines such as IL-12, TNF- α and IFN- γ ; and stimulate Th2 anti-inflammatory cytokine production such as IL-10, IL-4, and TGF- β [98]. Estradiol has also been shown to modulate the main activities (maturation, differentiation, and migration) of myeloid cells, including monocytes, macrophages, and dendritic cells [99-102]. Thus, estradiol has an

important impact on immune cells and affects both the innate and the adaptive immune systems, which may account for its contribution in diseases associated with immune disorder.

Estrogen and estrogen receptors in the gut

In an effort to identify extra-gonadal sites of *de novo* estradiol synthesis, we generated a double transgenic mouse line in which a transgenic aromatase (*cyp19*) promoter induces the expression of a red fluorescent protein (RFP) (un-published). In this animal, RFP signal is strongly expressed in the Peyer's patch (Pp), a secondary lymphoid organ in the intestine. Pp have an organizational structure similar to lymph nodes consisting of multiple follicles and interfollicular areas. A follicle is made of a germinal center that is filled with proliferating B-lymphocytes, follicular dendritic cells, and macrophages; the interfollicular area is populated with T-lymphocytes as well as B-lymphocytes, macrophages, and dendritic cells. As part of the gut-associated lymphoid tissue, Pp are known as inductive sites of intestinal immune responses [103]. The induction process in the Pp starts with sensing antigens or microbes in the gut lumen by M-cells located in a monolayer of specialized intestinal epithelial cells known as the follicle-associated epithelium. M-cells transport antigens to antigen-presenting cells, specifically dendritic cells (DCs), within the underlying sub-epithelial dome through transcytosis. Dendritic cells then further present antigens to T-and B-lymphocytes, triggering priming and proliferation of lymphocytes to complete the immune response. A well-known effect of the Pp's induction function is generating antigen-specific intestinal IgA responses, which is critical for maintaining host-microbiota interaction, generating immune tolerance, and preventing infection [104-106]. Interestingly, estrogens plays a significant role in the gastrointestinal tract. In this section, we will describe some of the lesser known roles for estrogen in the gastrointestinal system.

Napoleon Bonaparte was not aware of the true importance of his words when he said "*An army marches on its stomach.*" Technically, an army marches on its intestines. The gastrointestinal tract (GIT) is a unique environment colonized by a remarkable variety of bacteria as well as other organisms including fungi and viruses. This superorganism, the microbiome, is not a simple spectator in biological processes but is an active component of the biochemical and metabolic health of the host [107]. The microbiome is capable of digesting large molecules into simpler ones that can be efficiently reabsorbed by the host. The importance of a healthy microbiome has been well published [108-112], and multiple pathologies have been correlated with poor diversity of the microbiome, including irritable bowel (IBS)[113], osteoporosis [114, 115], and gluten intolerance [116]. Therefore, controlling the microbiome is paramount to maintaining an optimally functioning GIT. The mucosal epithelium is perfectly adapted to monitor both microbial and nutrient composition. The release of antimicrobial peptides [117] or anti-inflammatory molecules maintains the optimal microbial ecology depending on the current GIT contents.

Appetite. Researchers have noted a correlation between estradiol levels and appetite. Food intake is significantly decreased during the preovulatory period when estradiol levels are increasing [118]. These actions are attributed to estradiol inhibiting appetite indirectly through cannabinoid receptors [119]. Further,

blocking estrogen receptors with ICI182,270 ablates any action of estradiol on appetite [119]. What is more interesting is that appetite is influenced by the microbiome present in the GIT. Bacterial peptides signal hunger or satiation [117, 120]; in essence, the bacteria control our desire to eat. Locally synthesized estrogen produced in response to microbiome composition in turn may influence immune responses, bringing us back to control of microbiome composition.

Immune function. Estrogenic compounds in the gut lumen suppress immune function through targeted apoptosis and inhibition of cell proliferation in the germinal centers of ileal Pp [121]. The Pp are important in generating protective immune responses to pathogens through both innate and cell mediated responses [121] and are also key in tolerizing the host to food antigens. The mucosal surfaces of the gut must maintain homeostasis, allowing sufficient function of Pp to prevent immune responses to food antigens yet not responding prolifically to commensal bacteria in the gut. Abnormal Pp function through estrogenic compounds is responsible for initializing autoimmune responses and impaired innate responses. Again, we see the constituents of the gut signaling control of the microbiome composition. This leads into the next topic of estrogens and cancer.

Cancer. The small intestine is the main absorptive area of the gastrointestinal tract. To maximize absorption, the epithelial layer is covered with invaginations or crypts of Lieberkühn and exists as a sheet of single cells. These cells are prone to injury and are therefore replaced every 3-5 days [122]. To facilitate this replacement, the base of the crypts is populated with stem cells that differentiate into the mature epithelium as they migrate towards the crest of the crypt. ER α and ER β are both expressed in the crypt cells. However, they are distributed such that ER α is predominantly expressed in the cells at the base of crypts and ER β is expressed in the cells towards the crest. ER α signaling stimulates proliferation [123] and ER β signaling opposes this action [124, 125], and the net signaling from the two receptors controls proliferation. To further support the role of estrogen receptors in tumor development, ER β -deficient mice demonstrate a hyper-proliferation of the colonic epithelium with progression to colon carcinoma [91, 126]. More than 30 years ago, it was established that there is an associative risk between reduced estrogen levels and colorectal cancer in menopausal women [127] and that hormone (estrogen) replacement therapy reduces the incidence of colorectal cancer [128]. Recent literature on estrogen and colorectal cancer confirms an anti-tumorigenic role for estrogen signaling in the gut due to preferential ER β signaling [129].

However, estrogen in the gut is not always good. A recent review by Kwa *et al* [107] associated the “estrobolome” [130], bacteria with the capacity to metabolize estrogens, with level of risk for breast cancer. A phylogenetic diverse microbiome favors metabolism of conjugated estrogens. Once metabolized, the free estrogens are more easily reabsorbed increasing systemic estrogen levels. Increased circulating estrogens levels increases relative risk for hormone dependent malignancies such as breast cancer. As described above, our recent unpublished work has demonstrated that not only are Pp able to respond to estrogens, but they are also a significant site of estradiol synthesis. Thus, Pp are able to monitor the

bacterial diversity of the gut lumen and secrete estradiol. This estradiol then regulates immune responses locally and ultimately alters the diversity of the microbiome.

In conclusion, although estradiol is best recognized as sex hormone that regulates the development and function of reproductive hormone across the entire mammalian species, ever-growing evidence demonstrates its multi-faceted nature in exerting its role in non-reproductive organs and systems under normal as well as pathological conditions. It will be exciting to see what other functions estradiol may play in local tissues and from where the hormone is supplied to those sites.

References

1. Cutolo M, Capellino S, Sulli A et al (2006) Estrogens and autoimmune diseases. *Ann N Y Acad Sci* 1089, 538-547
2. Hogan, A.M., et al., Estrogen and gastrointestinal malignancy. *Mol Cell Endocrinol*, 307(1-2): p. 19-24
3. Hughes, G.C. and D. Choubey, Modulation of autoimmune rheumatic diseases by oestrogen and progesterone. *Nat Rev Rheumatol*, 2014. 10(12): p. 740-51
4. Brown, S.B. and S.E. Hankinson, Endogenous estrogens and the risk of breast, endometrial, and ovarian cancers. *Steroids*, 2014
5. Simpson, E., et al., Local estrogen biosynthesis in males and females. *Endocr Relat Cancer*, 1999. 6(2): p. 131-7
6. Simpson, E.R., Sources of estrogen and their importance. *J Steroid Biochem Mol Biol*, 2003. 86(3-5): p. 225-30
7. Nelson, L.R. and S.E. Bulun, Estrogen production and action. *J Am Acad Dermatol*, 2001. 45(3 Suppl): p. S116-24
8. Hess, R.A., et al., A role for oestrogens in the male reproductive system. *Nature*, 1997. 390(6659): p. 509-12
9. Hemsell, D.L., et al., Plasma precursors of estrogen. II. Correlation of the extent of conversion of plasma androstenedione to estrone with age. *J Clin Endocrinol Metab*, 1974. 38(3): p. 476-9
10. Labrie, F., et al., DHEA and the intracrine formation of androgens and estrogens in peripheral target tissues: its role during aging. *Steroids*, 1998. 63(5-6): p. 322-8
11. Ishunina, T.A., et al., Diminished aromatase immunoreactivity in the hypothalamus, but not in the basal forebrain nuclei in Alzheimer's disease. *Neurobiol Aging*, 2005. 26(2): p. 173-94
12. Azcoitia, I., et al., Aromatase expression by reactive astroglia is neuroprotective. *Ann N Y Acad Sci*, 2003. 1007: p. 298-305
13. Balthazart, J., et al., Distribution of aromatase-immunoreactive cells in the mouse forebrain. *Cell Tissue Res*, 1991. 263(1): p. 71-9
14. Yague, J.G., et al., Aromatase expression in the human temporal cortex. *Neuroscience*, 2006. 138(2): p. 389-401
15. Wehrenberg, U., J. Prange-Kiel, and G.M. Rune, Steroidogenic factor-1 expression in marmoset and rat hippocampus: co-localization with StAR and aromatase. *J Neurochem*, 2001. 76(6): p. 1879-86
16. Yague, J.G., et al., Aromatase distribution in the monkey temporal neocortex and hippocampus. *Brain Res*, 2008. 1209: p. 115-27
17. Cleland, W.H., C.R. Mendelson, and E.R. Simpson, Aromatase activity of membrane fractions of human adipose tissue stromal cells and adipocytes. *Endocrinology*, 1983. 113(6): p. 2155-60
18. Bulun, S.E., et al., A link between breast cancer and local estrogen biosynthesis suggested by quantification of breast adipose tissue aromatase cytochrome P450 transcripts using competitive

- polymerase chain reaction after reverse transcription. *J Clin Endocrinol Metab*, 1993. 77(6): p. 1622-8
19. Simpson, E.R., et al., Aromatase expression in health and disease. *Recent Prog Horm Res*, 1997. 52: p. 185-213; discussion 213-4
 20. Roselli, C.E. and J.A. Resko, Cytochrome P450 aromatase (CYP19) in the non-human primate brain: distribution, regulation, and functional significance. *J Steroid Biochem Mol Biol*, 2001. 79(1-5): p. 247-53
 21. Enjuanes, A., et al., Regulation of CYP19 gene expression in primary human osteoblasts: effects of vitamin D and other treatments. *Eur J Endocrinol*, 2003. 148(5): p. 519-26
 22. Watanabe, M., et al., Aromatase expression in the human fetal osteoblastic cell line SV-HFO. *J Mol Endocrinol*, 2004. 32(2): p. 533-45
 23. Janssen, J.M., et al., Estradiol formation by human osteoblasts via multiple pathways: relation with osteoblast function. *J Cell Biochem*, 1999. 75(3): p. 528-37
 24. Castagnetta, L.A., et al., Local estrogen formation by nontumoral, cirrhotic, and malignant human liver tissues and cells. *Cancer Res*, 2003. 63(16): p. 5041-5
 25. Hata, S., et al., Aromatase in human liver and its diseases. *Cancer Med*, 2013. 2(3): p. 305-15
 26. Robic, A., et al., Exploration of steroidogenesis-related genes in testes, ovaries, adrenals, liver and adipose tissue in pigs. *Anim Sci J*, 2016. 87(8): p. 1041-7
 27. Nicol, M.R., et al., Estrogen biosynthesis in human H295 adrenocortical carcinoma cells. *Mol Cell Endocrinol*, 2009. 300(1-2): p. 115-20
 28. Zhou, Y., et al., Ample Evidence: Dehydroepiandrosterone (DHEA) Conversion into Activated Steroid Hormones Occurs in Adrenal and Ovary in Female Rat. *PLoS One*, 2015. 10(5): p. e0124511
 29. Ueyama, T., et al., Gastric parietal cells: potent endocrine role in secreting estrogen as a possible regulator of gastro-hepatic axis. *Endocrinology*, 2002. 143(8): p. 3162-70
 30. Leshin, M., et al., Increased estrogen formation and aromatase activity in fibroblasts cultured from the skin of chickens with the Henny feathering trait. *J Biol Chem*, 1981. 256(9): p. 4341-4
 31. Pomari, E., et al., Intracrine sex steroid synthesis and signaling in human epidermal keratinocytes and dermal fibroblasts. *FASEB J*, 2015. 29(2): p. 508-24
 32. Berkovitz, G.D., T.R. Brown, and M. Fujimoto, Aromatase activity in human skin fibroblasts grown in cell culture. *Steroids*, 1987. 50(1-3): p. 281-95
 33. Harada, N., et al., Localized expression of aromatase in human vascular tissues. *Circ Res*, 1999. 84(11): p. 1285-91
 34. Bayard, F., et al., Estrogen synthesis, estrogen metabolism, and functional estrogen receptors in rat arterial smooth muscle cells in culture. *Endocrinology*, 1995. 136(4): p. 1523-9
 35. Bayard, F., et al., Oestrogen synthesis, oestrogen metabolism and functional oestrogen receptors in bovine aortic endothelial cells. *Ciba Found Symp*, 1995. 191: p. 122-32; discussion 132-8
 36. Samy, T.S., et al., Divergent immune responses in male and female mice after trauma-hemorrhage: dimorphic alterations in T lymphocyte steroidogenic enzyme activities. *Endocrinology*, 2001. 142(8): p. 3519-29
 37. Abraham, I.M., et al., Estrogen receptor beta mediates rapid estrogen actions on gonadotropin-releasing hormone neurons in vivo. *J Neurosci*, 2003. 23(13): p. 5771-7
 38. Higaki, S., et al., Response of ERbeta and aromatase expression in the monkey hippocampal formation to ovariectomy and menopause. *Neurosci Res*, 2012. 72(2): p. 148-54
 39. Shughrue, P.J., P.J. Scrimo, and I. Merchenthaler, Estrogen binding and estrogen receptor characterization (ERalpha and ERbeta) in the cholinergic neurons of the rat basal forebrain. *Neuroscience*, 2000. 96(1): p. 41-9
 40. Almey, A., T.A. Milner, and W.G. Brake, Estrogen receptor alpha and G-protein coupled estrogen receptor 1 are localized to GABAergic neurons in the dorsal striatum. *Neurosci Lett*, 2016. 622: p. 118-23
 41. Xu, Y., et al., Distinct hypothalamic neurons mediate estrogenic effects on energy homeostasis and reproduction. *Cell Metab*, 2011. 14(4): p. 453-65
 42. Cooke, P.S. and A. Naaz, Role of estrogens in adipocyte development and function. *Exp Biol Med* (Maywood), 2004. 229(11): p. 1127-35

43. Davis, K.E., et al., Sexually dimorphic role of G protein-coupled estrogen receptor (GPER) in modulating energy homeostasis. *Horm Behav*, 2014. 66(1): p. 196-207
44. Dieudonne, M.N., et al., Evidence for functional estrogen receptors alpha and beta in human adipose cells: regional specificities and regulation by estrogens. *Am J Physiol Cell Physiol*, 2004. 286(3): p. C655-61
45. Sjogren, K., et al., Elevated aromatase expression in osteoblasts leads to increased bone mass without systemic adverse effects. *J Bone Miner Res*, 2009. 24(7): p. 1263-70
46. Martin-Millan, M., et al., The estrogen receptor-alpha in osteoclasts mediates the protective effects of estrogens on cancellous but not cortical bone. *Mol Endocrinol*, 2010. 24(2): p. 323-34
47. Maatta, J.A., et al., Inactivation of estrogen receptor alpha in bone-forming cells induces bone loss in female mice. *FASEB J*, 2013. 27(2): p. 478-88
48. Crusode de Souza, M., E. Sasso-Cerri, and P.S. Cerri, Immunohistochemical detection of estrogen receptor beta in alveolar bone cells of estradiol-treated female rats: possible direct action of estrogen on osteoclast life span. *J Anat*, 2009. 215(6): p. 673-81
49. Braidman, I.P., et al., Localization of estrogen receptor beta protein expression in adult human bone. *J Bone Miner Res*, 2001. 16(2): p. 214-20
50. Bord, S., et al., Estrogen receptors alpha and beta are differentially expressed in developing human bone. *J Clin Endocrinol Metab*, 2001. 86(5): p. 2309-14
51. Karas, R.H., B.L. Patterson, and M.E. Mendelsohn, Human vascular smooth muscle cells contain functional estrogen receptor. *Circulation*, 1994. 89(5): p. 1943-50
52. Russell, K.S., et al., Human vascular endothelial cells contain membrane binding sites for estradiol, which mediate rapid intracellular signaling. *Proc Natl Acad Sci U S A*, 2000. 97(11): p. 5930-5
53. Broughton, B.R., A.A. Miller, and C.G. Sobey, Endothelium-dependent relaxation by G protein-coupled receptor 30 agonists in rat carotid arteries. *Am J Physiol Heart Circ Physiol*, 2010. 298(3): p. H1055-61
54. Chakrabarti, S. and S.T. Davidge, High glucose-induced oxidative stress alters estrogen effects on ERalpha and ERbeta in human endothelial cells: reversal by AMPK activator. *J Steroid Biochem Mol Biol*, 2009. 117(4-5): p. 99-106
55. Campbell-Thompson, M., K.K. Reyher, and L.B. Wilkinson, Immunolocalization of estrogen receptor alpha and beta in gastric epithelium and enteric neurons. *J Endocrinol*, 2001. 171(1): p. 65-73
56. Gaudet, H.M., et al., The G-protein coupled estrogen receptor, GPER: The inside and inside-out story. *Mol Cell Endocrinol*, 2015. 418 Pt 3: p. 207-19
57. Thornton, M.J., et al., Oestrogen receptor beta is the predominant oestrogen receptor in human scalp skin. *Exp Dermatol*, 2003. 12(2): p. 181-90
58. El Safoury, O., L. Rashid, and M. Ibrahim, A study of androgen and estrogen receptors alpha, beta in skin tags. *Indian J Dermatol*, 2010. 55(1): p. 20-4
59. Cesinaro, A.M., L. Roncati, and A. Maiorana, Estrogen receptor alpha overexpression in multinucleate cell angiohistiocytoma: new insights into the pathogenesis of a reactive process. *Am J Dermatopathol*, 2010. 32(7): p. 655-9
60. Trejter, M., et al., Expression of estrogen, estrogen related and androgen receptors in adrenal cortex of intact adult male and female rats. *Folia Histochem Cytobiol*, 2015. 53(2): p. 133-44
61. Velders, M., et al., Selective estrogen receptor-beta activation stimulates skeletal muscle growth and regeneration. *FASEB J*, 2012. 26(5): p. 1909-20
62. Potier, M., et al., Expression and regulation of estrogen receptors in mesangial cells: influence on matrix metalloproteinase-9. *J Am Soc Nephrol*, 2001. 12(2): p. 241-51
63. Le May, C., et al., Estrogens protect pancreatic beta-cells from apoptosis and prevent insulin-deficient diabetes mellitus in mice. *Proc Natl Acad Sci U S A*, 2006. 103(24): p. 9232-7
64. Liu, S., et al., Importance of extranuclear estrogen receptor-alpha and membrane G protein-coupled estrogen receptor in pancreatic islet survival. *Diabetes*, 2009. 58(10): p. 2292-302
65. Gao, H., et al., Long-term administration of estradiol decreases expression of hepatic lipogenic genes and improves insulin sensitivity in ob/ob mice: a possible mechanism is through direct regulation of signal transducer and activator of transcription 3. *Mol Endocrinol*, 2006. 20(6): p. 1287-99

66. Reue, K. and J.R. Dwyer, Lipin proteins and metabolic homeostasis. *J Lipid Res*, 2009. 50 Suppl: p. S109-14
67. Miller, W.R., Aromatase activity in breast tissue. *J Steroid Biochem Mol Biol*, 1991. 39(5B): p. 783-90
68. Sasano, H., et al., Aromatase in human bone tissue. *J Bone Miner Res*, 1997. 12(9): p. 1416-23
69. Shozu, M. and E.R. Simpson, Aromatase expression of human osteoblast-like cells. *Mol Cell Endocrinol*, 1998. 139(1-2): p. 117-29
70. Zhong, M., et al., 17beta-Estradiol regulates rat growth plate chondrocyte apoptosis through a mitochondrial pathway not involving nitric oxide or MAPKs. *Endocrinology*, 2011. 152(1): p. 82-92
71. Slominski, A. and J. Wortsman, Neuroendocrinology of the skin. *Endocr Rev*, 2000. 21(5): p. 457-87
72. Brincat, M.P., Hormone replacement therapy and the skin. *Maturitas*, 2000. 35(2): p. 107-17
73. Harada, N., A unique aromatase (P-450AROM) mRNA formed by alternative use of tissue-specific exons 1 in human skin fibroblasts. *Biochem Biophys Res Commun*, 1992. 189(2): p. 1001-7
74. Emoto, N., N. Ling, and A. Baird, Growth factor-mediated regulation of aromatase activity in human skin fibroblasts. *Proc Soc Exp Biol Med*, 1991. 196(3): p. 351-8
75. Barros, R.P. and J.A. Gustafsson, Estrogen receptors and the metabolic network. *Cell Metab*, 2011. 14(3): p. 289-99
76. Parthasarathy, C., V.N. Renuka, and K. Balasubramanian, Sex steroids enhance insulin receptors and glucose oxidation in Chang liver cells. *Clin Chim Acta*, 2009. 399(1-2): p. 49-53
77. Klair, J.S., et al., A longer duration of estrogen deficiency increases fibrosis risk among postmenopausal women with nonalcoholic fatty liver disease. *Hepatology*, 2016. 64(1): p. 85-91.
78. Iavarone, M., et al., The clinical and pathogenetic significance of estrogen receptor-beta expression in chronic liver diseases and liver carcinoma. *Cancer*, 2003. 98(3): p. 529-34
79. Marzioni, M., et al., An oestrogen receptor beta-selective agonist exerts anti-neoplastic effects in experimental intrahepatic cholangiocarcinoma. *Dig Liver Dis*, 2012. 44(2): p. 134-42
80. Colciago, A., et al., Dimorphic expression of testosterone metabolizing enzymes in the hypothalamic area of developing rats. *Brain Res Dev Brain Res*, 2005. 155(2): p. 107-16
81. Rettberg, J.R., J. Yao, and R.D. Brinton, Estrogen: a master regulator of bioenergetic systems in the brain and body. *Front Neuroendocrinol*, 2014. 35(1): p. 8-30
82. Kaludjerovic, J. and W.E. Ward, The Interplay between Estrogen and Fetal Adrenal Cortex. *J Nutr Metab*, 2012. 2012: p. 837901
83. Alonso-Magdalena, P., et al., Pancreatic insulin content regulation by the estrogen receptor ER alpha. *PLoS One*, 2008. 3(4): p. e2069
84. Yuchi, Y., et al., Estrogen Receptor alpha Regulates beta-Cell Formation During Pancreas Development and Following Injury. *Diabetes*, 2015. 64(9): p. 3218-28
85. Pedram, A., et al., Estrogen receptor-beta prevents cardiac fibrosis. *Mol Endocrinol*, 2010. 24(11): p. 2152-65
86. Novensa, L., et al., Aging negatively affects estrogens-mediated effects on nitric oxide bioavailability by shifting ERalpha/ERbeta balance in female mice. *PLoS One*, 2011. 6(9): p. e25335
87. Prossnitz, E.R. and M. Maggiolini, Mechanisms of estrogen signaling and gene expression via GPR30. *Mol Cell Endocrinol*, 2009. 308(1-2): p. 32-8
88. Lowe, D.A., K.A. Baltgalvis, and S.M. Greising, Mechanisms behind estrogen's beneficial effect on muscle strength in females. *Exerc Sport Sci Rev*, 2010. 38(2): p. 61-7
89. Enns, D.L. and P.M. Tiidus, The influence of estrogen on skeletal muscle: sex matters. *Sports Med*, 2010. 40(1): p. 41-58
90. Antus, B., et al., Estradiol is nephroprotective in the rat remnant kidney. *Nephrol Dial Transplant*, 2003. 18(1): p. 54-61
91. Wada-Hiraike, O., et al., Role of estrogen receptor beta in colonic epithelium. *Proc Natl Acad Sci U S A*, 2006. 103(8): p. 2959-64

92. Shen, L. and J.R. Turner, Role of epithelial cells in initiation and propagation of intestinal inflammation. Eliminating the static: tight junction dynamics exposed. *Am J Physiol Gastrointest Liver Physiol*, 2006. 290(4): p. G577-82
93. Straub, R.H., The complex role of estrogens in inflammation. *Endocr Rev*, 2007. 28(5): p. 521-74
94. Phiel, K.L., et al., Differential estrogen receptor gene expression in human peripheral blood mononuclear cell populations. *Immunol Lett*, 2005. 97(1): p. 107-13
95. Kouro, T., et al., Characteristics of early murine B-lymphocyte precursors and their direct sensitivity to negative regulators. *Blood*, 2001. 97(9): p. 2708-15
96. Masuzawa, T., et al., Estrogen deficiency stimulates B lymphopoiesis in mouse bone marrow. *J Clin Invest*, 1994. 94(3): p. 1090-7
97. Wilson, C.A., S.A. Mrose, and D.W. Thomas, Enhanced production of B lymphocytes after castration. *Blood*, 1995. 85(6): p. 1535-9
98. Salem, M.L., Estrogen, a double-edged sword: modulation of TH1- and TH2-mediated inflammations by differential regulation of TH1/TH2 cytokine production. *Curr Drug Targets Inflamm Allergy*, 2004. 3(1): p. 97-104
99. Harkonen, P.L. and H.K. Vaananen, Monocyte-macrophage system as a target for estrogen and selective estrogen receptor modulators. *Ann N Y Acad Sci*, 2006. 1089: p. 218-27
100. Komi, J. and O. Lassila, Nonsteroidal anti-estrogens inhibit the functional differentiation of human monocyte-derived dendritic cells. *Blood*, 2000. 95(9): p. 2875-82
101. Mao, A., et al., Estrogen selectively promotes the differentiation of dendritic cells with characteristics of Langerhans cells. *J Immunol*, 2005. 175(8): p. 5146-51
102. Mor, G., et al., Interaction of the estrogen receptors with the Fas ligand promoter in human monocytes. *J Immunol*, 2003. 170(1): p. 114-22
103. Jung, C., J.P. Hugot, and F. Barreau, Peyer's Patches: The Immune Sensors of the Intestine. *Int J Inflam*, 2010. 2010: p. 823710
104. Lycke, N.Y. and M. Bemark, The role of Peyer's patches in synchronizing gut IgA responses. *Front Immunol*, 2012. 3: p. 329
105. Mkaddem, S.B., et al., IgA, IgA receptors, and their anti-inflammatory properties. *Curr Top Microbiol Immunol*, 2014. 382: p. 221-35
106. Singh, K., C. Chang, and M.E. Gershwin, IgA deficiency and autoimmunity. *Autoimmun Rev*, 2014. 13(2): p. 163-77
107. Kwa, M., et al., The Intestinal Microbiome and Estrogen Receptor-Positive Female Breast Cancer. *J Natl Cancer Inst*, 2016. 108(8)
108. Relman, D.A., Gut microbiota: How to build healthy growth-promoting gut communities. *Nat Rev Gastroenterol Hepatol*, 2016. 13(7): p. 379-80
109. Wischmeyer, P.E., D. McDonald, and R. Knight, Role of the microbiome, probiotics, and 'dysbiosis therapy' in critical illness. *Curr Opin Crit Care*, 2016. 22(4): p. 347-53
110. Nishijima, S., et al., The gut microbiome of healthy Japanese and its microbial and functional uniqueness. *DNA Res*, 2016. 23(2): p. 125-33
111. Highet, A.R., et al., Gut microbiome in sudden infant death syndrome (SIDS) differs from that in healthy comparison babies and offers an explanation for the risk factor of prone position. *Int J Med Microbiol*, 2014. 304(5-6): p. 735-41
112. Tuddenham, S. and C.L. Sears, The intestinal microbiome and health. *Curr Opin Infect Dis*, 2015. 28(5): p. 464-70
113. Fourie, N.H., et al., The microbiome of the oral mucosa in irritable bowel syndrome. *Gut Microbes*, 2016: p. 1-16
114. Hernandez, C.J., et al., Links Between the Microbiome and Bone. *J Bone Miner Res*, 2016
115. Weaver, C.M., Diet, gut microbiome, and bone health. *Curr Osteoporos Rep*, 2015. 13(2): p. 125-30
116. Fernandez-Feo, M., et al., The cultivable human oral gluten-degrading microbiome and its potential implications in coeliac disease and gluten sensitivity. *Clin Microbiol Infect*, 2013. 19(9): p. E386-94

117. Zhao, Z. and T. Sakai, Characteristic features of ghrelin cells in the gastrointestinal tract and the regulation of stomach ghrelin expression and production. *World J Gastroenterol*, 2008. 14(41): p. 6306-11
118. Lyons, P.M., et al., Reduction of food intake in the ovulatory phase of the menstrual cycle. *Am J Clin Nutr*, 1989. 49(6): p. 1164-8
119. Mela, V., et al., Modulatory influences of estradiol and other anorexigenic hormones on metabotropic, Gi/o-coupled receptor function in the hypothalamic control of energy homeostasis. *J Steroid Biochem Mol Biol*, 2016. 160: p. 15-26
120. Blaut, M., Gut microbiota and energy balance: role in obesity. *Proc Nutr Soc*, 2015. 74(3): p. 227-34
121. Obremski, K. and G. Poniatowska-Broniek, Zearalenone induces apoptosis and inhibits proliferation in porcine ileal Peyer's patch lymphocytes. *Pol J Vet Sci*, 2015. 18(1): p. 153-61
122. Matthews, J., et al., Estrogen receptor (ER) beta modulates ERalpha-mediated transcriptional activation by altering the recruitment of c-Fos and c-Jun to estrogen-responsive promoters. *Mol Endocrinol*, 2006. 20(3): p. 534-43
123. Matthews, J. and J.A. Gustafsson, Estrogen signaling: a subtle balance between ER alpha and ER beta. *Mol Interv*, 2003. 3(5): p. 281-92
124. Matthews, J. and J.A. Gustafsson, Estrogen receptor and aryl hydrocarbon receptor signaling pathways. *Nucl Recept Signal*, 2006. 4: p. e016
125. Liu, M.M., et al., Opposing action of estrogen receptors alpha and beta on cyclin D1 gene expression. *J Biol Chem*, 2002. 277(27): p. 24353-60
126. D'Errico, I. and A. Moschetta, Nuclear receptors, intestinal architecture and colon cancer: an intriguing link. *Cell Mol Life Sci*, 2008. 65(10): p. 1523-43
127. Butcher, D., et al., Female gender is a major determinant of changing subsite distribution of colorectal cancer with age. *Cancer*, 1985. 56(3): p. 714-6
128. Chlebowski, R.T., et al., Estrogen plus progestin and colorectal cancer in postmenopausal women. *N Engl J Med*, 2004. 350(10): p. 991-1004
129. Caiazza, F., et al., Estrogen receptors and their implications in colorectal carcinogenesis. *Front Oncol*, 2015. 5: p. 19
130. Plottel, C.S. and M.J. Blaser, Microbiome and malignancy. *Cell Host Microbe*, 2011. 10(4): p. 324-35