

BMB Reports – Manuscript Submission

Manuscript Draft

**Manuscript Number:** BMB-20-179

**Title:** Activation of pannexin-1 mediates triglyceride-induced macrophage cell death

**Article Type:** Article

**Keywords:** Triglyceride (TG); Potassium efflux; Pannexin-1; THP-1 macrophages; Cell death

**Corresponding Author:** Yoon Suk Kim

**Authors:** Byung Chul Jung<sup>1,2</sup>, Sung Hoon Kim<sup>2,3</sup>, Jaewon Lim<sup>2,4</sup>, Yoon Suk Kim<sup>2,\*</sup>

**Institution:** <sup>1</sup>Department of Nutritional Sciences and Toxicology, University of California, Berkeley, State of California 94720, United States,

<sup>2</sup>Department of Biomedical Laboratory Science, College of Health Sciences, Yonsei University, Wonju, Gangwon-do 26493, Republic of Korea,

<sup>3</sup>Department of Biomedical Laboratory Science, Korea Nazarene University, Cheonan, Chungcheongnam-do 31172, Republic of Korea,

<sup>4</sup>Department of Biomedical Laboratory Science, College of Medical Sciences, Daegu Haany University, Gyeongsan, Gyeongsangbuk-do 38610, Republic of Korea,

**Manuscript Type:** Article

**Title:** Activation of pannexin-1 mediates triglyceride-induced macrophage cell death

**Author's name:** Byung Chul Jung<sup>1,2</sup>, Sung Hoon Kim<sup>2,3</sup>, Jaewon Lim<sup>2,4,\*</sup>, and Yoon Suk Kim<sup>2,\*</sup>

**Affiliation:** <sup>1</sup> Department of Nutritional Sciences and Toxicology, University of California, Berkeley, State of California 94720, United States

<sup>2</sup> Department of Biomedical Laboratory Science, College of Health Sciences, Yonsei University, Wonju, Gangwon-do 26493, Republic of Korea

<sup>3</sup> Department of Biomedical Laboratory Science, Korea Nazarene University, Cheonan, Chungcheongnam-do 31172, Republic of Korea

<sup>4</sup> Department of Biomedical Laboratory Science, College of Medical Sciences, Daegu Haany University, Gyeongsan, Gyeongsangbuk-do 38610, Republic of Korea

**Running Title:** Pannexin-1 in TG-induced macrophage cell death

**Keywords:** Triglyceride (TG), Potassium efflux, Pannexin-1, THP-1 macrophages, Cell death

**Corresponding Author's Information:** \*To whom correspondence should be addressed:

Yoon Suk Kim, Department of Biomedical Laboratory Science, College of Health Science,  
Yonsei University, Wonju, Gangwon-do, 26493, Republic of Korea

Tel : 82-33-760-2860, E-mail : yoonsukkim@yonsei.ac.kr

Jaewon Lim, Department of Biomedical Laboratory Science, College of Medical Sciences,  
Daegu Haany University, Gyeongsan, Gyeongsangbuk-do 38610, Republic of Korea

Tel : 82-53-819-1352, E-mail : jaewon330@dhu.ac.kr

**ABSTRACT**

The accumulation of triglycerides (TGs) in macrophages induces cell death, a risk factor in the pathogenesis of atherosclerosis. We had previously reported that TG-induced macrophage death is triggered by caspase-1 and -2, therefore we investigated the mechanism underlying this phenomenon. We found that potassium efflux is increased in TG-treated THP-1 macrophages and that the inhibition of potassium efflux blocks TG-induced cell death as well as caspase-1 and -2 activation. Furthermore, reducing ATP concentration (known to induce potassium efflux), restored cell viability and caspase-1 and -2 activity. The activation of pannexin-1 (a channel that releases ATP), was increased after TG treatment in THP-1 macrophages. Inhibition of pannexin-1 activity using its inhibitor, probenecid, recovered cell viability and blocked the activation of caspase-1 and -2 in TG-treated macrophages. These results suggest that TG-induced THP-1 macrophage cell death is induced via pannexin-1 activation, which increases extracellular ATP, leading to an increase in potassium efflux.

## INTRODUCTION

Atherosclerosis is a disease characterized by the narrowing and hardening of arteries due to the formation of lipid-laden plaques, often leading to ischemic cardiomyopathy and stroke (1). Since the abnormal accumulation of lipoproteins, especially oxidative low-density lipoproteins (Oxi-LDL), is considered the first step in the pathogenesis of atherosclerosis, many studies have been directed towards understanding the characteristics of Oxi-LDL and oxidative stress (2, 3). Multiple lines of evidence support that elevated TG levels are associated with an increased risk of atherosclerosis and coronary heart disease (4, 5). We have previously demonstrated that TGs induced the macrophage death via caspase-1 and 2 (6, 7). These studies suggested a putative mechanism by which TG contributes to the development of atherosclerosis. However, the mechanism by which TG triggers the caspase cascade in macrophages remains elusive.

Potassium efflux occurs during cell death due to loss of ionic homeostasis. Furthermore, dysregulation of potassium efflux is one of the causative factors of caspase-dependent cell death (8). For example, the potassium ionophore valinomycin can induce potassium efflux, which triggers caspase-3 activation and subsequent cell death in CHO cells (9). Furthermore,  $\alpha$ -toxin-mediated caspase-2 activation and cell death were suppressed on preventing potassium efflux in HeLa cells, indicating that potassium efflux is a prerequisite for caspase-2-mediated cell death (10). Many studies have speculated on the mechanisms by which potassium efflux is regulated. One of the well-known mechanism for increasing potassium efflux is the opening of the ATP-sensitive potassium channel (11). It has also been reported that ATP-sensitive potassium channels open during the initial phase of myocardial ischemia, one of the consequences of atherosclerosis (12).

Pannexins are large pore-forming channels and consist of three family members (pannexin-1, 2, and 3). Since pannexin-1 is ubiquitously expressed ATP-permeable channel

and is involved in a variety of cellular processes, including cell death, it has attracted significant attention in research (13). In the current study, we investigated the involvement of potassium efflux in TG-induced macrophage cell death. We report that TG induces an increase of extracellular ATP via pannexin-1, which in turn activates ATP-sensitive potassium channels, resulting to an increase in potassium efflux and a subsequent activation of the caspase cascade in THP-1 macrophages. These results suggest that TG-induced cell death of macrophages is mediated by the activation of pannexin-1.

## RESULTS

### Potassium efflux is involved in TG-induced THP-1 macrophage cell death

Though previous studies have reported that TG-induced cell death is mediated by caspase-1 and 2 (6, 7), the mechanism by which they are activated is not fully understood. Several studies have reported that the release of intracellular potassium ions activates caspases (14, 15). Therefore, this study investigated whether potassium efflux is involved in TG-induced caspase-mediated THP-1 macrophage death. When extracellular potassium concentration was measured in PMA-differentiated THP-1 macrophages (incubated with or without 1 mg/mL of TG for 24 h), it was found to be higher in TG-treated macrophages (Fig. 1A). Furthermore, when macrophages were treated with glyburide, an inhibitor of ATP-sensitive potassium channels, TG-induced cell death was restored in a dose-dependent manner (Fig. 1B). When the extracellular concentration of potassium was increased via treatment with 25 mM KCl to prevent the release of potassium ions, cell viability was recovered (Fig. 1C). Suppression of potassium efflux by treating macrophages with glyburide reduced the levels of cleaved PARP and several caspases (caspase-3, 7, 8, and 9) that are known to be responsible for TG-induced macrophage death (Fig. 1D). When THP-1 macrophages were incubated with TG in the presence of glyburide, treatment with glyburide decreased the activity of caspase-1 which is reported to induce the activity of caspase-3, -7, -8, and -9 (7) (Fig. 1E and 1F). Treatment with glyburide also reduced the activity of caspase-2, a known upstream molecule of caspase-1 (7), which increased upon TG treatment (Fig. 1G). These results demonstrate that potassium efflux induces TG-stimulated macrophage death via the activation of caspases.

### The increase in extracellular ATP leads to TG-triggered macrophage cell death

It has recently been reported that an increase in extracellular ATP affects ATP-sensitive potassium channels, thereby increasing potassium efflux (16). Therefore, to

determine if extracellular ATP is involved in TG-induced apoptotic cell death, PMA-differentiated THP-1 macrophages were incubated with TG in the presence or absence of ATP to assess cell viability. Treatment of THP-1 macrophages with ATP and TG resulted in a greater reduction in cell viability compared to treatment with TG only (Fig. 2A). Subsequently, to investigate whether extracellular ATP enhances apoptotic cell death in TG-treated macrophages, PARP cleavage was confirmed by western blot analysis. Extracellular ATP increased the levels of cleaved PARP in TG-treated macrophages (Fig. 2B). In addition, extracellular ATP also increased the activity of caspase-1 and 2 in TG-stimulated macrophages (Fig. 2C and 2D). To elucidate whether the increase in extracellular ATP was responsible for an increased TG-triggered apoptotic cell death, THP-1 macrophages were treated with TG in the presence of the ATP-depleting enzyme apyrase, which catalyzes the hydrolysis of ATP and ADP to AMP and inorganic phosphate. Cell viability was recovered (Fig. 2E) and the cleavage of PARP and apoptotic caspases (caspase-3, -7, -8, and -9) was reduced in the presence of apyrase in TG-treated THP-1 macrophages (Fig. 2F). Moreover, ATP depletion decreased the activity of caspase-1 and 2, which was increased in response to TG treatment (Fig. 2G and 2H). These results suggested that extracellular ATP is associated with TG-triggered macrophage death.

### **Activation of the pannexin-1 channel mediates TG-induced THP-1 macrophage cell death**

Pannexin-1 channels are mechanosensitive conduits for ATP release in human cells (17). These channels are activated upon the cleavage of their C-terminus, and as they are activated, their cleavage fragments accumulate in the intracellular pool (18). To elucidate whether pannexin-1 is involved in macrophage cell death following TG treatment, THP-1 macrophages were treated with TG, and cytoplasmic pannexin-1 cleavage was measured using Western blot analysis. As a result, TG treatment increased cleaved pannexin-1 in a dose-



dependent manner and pannexin-1 cleavage began to increase from 1.5 h after treatment (Fig. 3A).

When the pannexin-1 channel is activated, both ATP and small dye molecules can pass through it (18). An Et-Br uptake assay was performed to evaluate the activation of the pannexin-1 channel after THP-1 macrophages were treated with TG. More fluorescent red dots were observed in TG-treated macrophages than those that were not (Fig. 3B and 3C). These results show that TG treatment increases pannexin-1 channel activity in THP-1 macrophages.

To determine whether the activity of pannexin-1 is involved in TG-induced macrophage death, TG-treated THP-1 macrophages were treated with probenecid, an inhibitor of pannexin-1. When the activity of pannexin-1 in TG-treated THP-1 macrophages was inhibited by probenecid, cell viability was recovered (Fig. 3D), and the cleavage of caspase-3, -7, -8, and -9, and PARP were decreased (Fig. 3E). The activity of caspase-1 and caspase-2 in TG-treated THP-1 macrophages were also reduced (Fig. 3F and 3G). Taken together, these results indicate that pannexin-1 channel activation mediates TG-triggered cell death in THP-1 macrophages.

### **TG-triggered cell death is mediated by ATP released by pannexin-1 channel activation**

To test whether the activation of the pannexin-1 channel contributes to extracellular ATP-dependent caspase-1 and -2 activation and increases TG-induced cell death, THP-1 macrophages were incubated with TG in the presence of probenecid and ATP. The cell viability recovered upon the inhibition of the pannexin-1 channels was reduced again by the presence of the extracellular ATP in a dose-dependent manner (Fig. 4A). In addition, the cleavage of the caspases and PARP was increased again in response to the presence of ATP (Fig. 4B). The activity of caspase-1 and 2 decreased upon probenecid treatment was increased again in response to ATP treatment in a dose-dependent manner (Fig. 4C and 4D). Taken together, these

results suggest that the activation of the pannexin-1 channel mediates ATP-dependent TG-triggered cell death in THP-1 macrophages.

## DISCUSSION

Macrophage cell death is an essential phenomenon in atherosclerosis, contributing to the development of plaques in advanced lesions (19). Since Oxi-LDL has been considered a key player in the development of atherosclerosis, much research has been directed at understanding the association between Oxi-LDL and macrophage death (20, 21). In contrast to Oxi-LDL, the role of TG in atherosclerosis has long been controversial. However, emerging evidence, including a recent large cohort study, suggested that elevated TG levels can be a causal risk factor for atherosclerosis (22). Thus, in the current study, we showed that i) TG increased extracellular ATP via pannexin-1, which in turn, ii) activates ATP-sensitive potassium channels, and that iii) there is an activation of the caspase cascade through increased potassium efflux. These results suggest that the TG-induced cell death of macrophages is mediated by pannexin-1 activation.

Pannexin-1 was identified as a plasma membrane hemichannel that releases ATP from the cytosol to the extracellular space during apoptosis (23). The released ATP can act as a “find-me” signal to recruit phagocytes and activate the initiation of inflammation, which can be an important contributor for the pathogenesis of atherosclerosis (23). We previously demonstrated that TG treatment enhanced the secretion of a pro-inflammatory cytokine, IL-1 $\beta$ , in THP-1 macrophages (6). Another function of increased extracellular ATP is the activation of ATP-sensitive potassium channels, subsequently leading to an increase in potassium efflux. Several investigations suggested that potassium channel activation contributes to the pathogenesis of atherosclerosis (24, 25). Ling and colleagues showed that the inhibition of ATP-sensitive potassium channels ameliorated atherosclerosis in ApoE<sup>-/-</sup> mouse models, directly elucidating the role of ATP-sensitive potassium channels in atherosclerosis (25). Moreover, they observed that the number of ATP-sensitive potassium channels in macrophages increased in atherosclerotic plaques, which may be explained by the present study. Here, we

showed that TG-mediated macrophage cell death was dependent on the activation of ATP-sensitive potassium channels. It is possible that increased ATP-sensitive potassium channels in macrophages cause an increase in potassium efflux, leading to macrophage cell death. This may then potentiate the development of more vulnerable plaques.

In conclusion, our results provide a novel mechanism regarding how TG induces macrophage cell death. Since an enhanced level of macrophage death is considered a crucial phenomenon in plaque development in atherosclerosis, our results imply that TG-induced macrophages can create plaques and produce thrombi, resulting in clinical ischemic cardiomyopathy. Furthermore, our results support that TG can be a risk factor in the pathogenesis of atherosclerosis, and that control of TG levels can be a valuable therapeutic target for the disease.

## MATERIALS AND METHODS

### Materials

Lipofundin<sup>®</sup> MCT/LCT 20% (B. Braun Melsungen AG; Melsungen, Germany) was used to deliver TG into cells, as previously described (26). Hereafter, Lipofundin<sup>®</sup> MCT/LCT 20% will be referred to as TG for convenience. The caspase-1 substrate, Ac-YVAD-pNA was purchased from Biomol (Plymouth Meeting, PA, USA). The caspase-2 substrate, Ac-VAIVAD-pNA was purchased from Sigma-Aldrich (St. Louis, MO, USA). Glyburide, apyrase, and probenecid were purchased from Sigma-Aldrich. The caspase-1 specific inhibitor, z-YVAD-FMK, and caspase-2 specific inhibitor, z-VDVAD-FMK, were purchased from BioVision (Mountain View, CA, USA). ATP was purchased from Sigma-Aldrich. Antibodies specific for caspase-1, -3, -7, -8, and -9, as well as PARP, were purchased from Cell Signaling Technology (Danvers, MA, USA). Antibodies specific for pannexin-1 were purchased from R&D systems (Minneapolis, MN, USA).

### Cell culture

The THP-1 human acute monocytic leukemia cell line (ATCC, Manassas, VA, USA) was grown in RPMI 1640 media supplemented with 10% fetal bovine serum and penicillin-streptomycin and maintained at 37°C in a humidified atmosphere with 5% CO<sub>2</sub>. To induce differentiation of THP-1 cells into macrophages, cells were seeded in 6-well plates at a density of  $1 \times 10^6$  cells/well and treated with 200 nM phorbol 12-myristate 13-acetate (PMA) for 48 h.

### Atomic absorption spectroscopy for measurement of K<sup>+</sup> concentration

To assess the concentration of K<sup>+</sup> efflux, THP-1 macrophages ( $1 \times 10^6$  cells per well) were incubated with or without TG for 24 h. Cells were washed with PBS and cultured in serum-free medium another 6 h at 37°C. The concentration of K<sup>+</sup> in each sample was quantified

using an atomic absorption spectrometer (AA7000F, Shimadzu, Tokyo, Japan).

### **Trypan blue dye exclusion assay**

To enumerate viable cells, cells were trypsinized and 10  $\mu$ L of 0.4% trypan blue stain solution was mixed with 10  $\mu$ L of the trypsinized cell suspension. Non-stained cells in the resulting mixture were counted using a hemocytometer (Marienfeld, Lauda-Königshofen, Germany).

### **Measurement of caspase activity**

The activities of caspase-1 and caspase-2 were determined as previously described (27). Briefly, cells were lysed with PBS buffer containing 1% Triton X-100 and centrifuged at  $19,000 \times g$  for 10 min at 4°C. The supernatant was collected and the total protein concentration was quantified. To detect caspase-1 activity, 90  $\mu$ g of the protein sample was combined with 200  $\mu$ M of Ac-YVAD-pNA in 150  $\mu$ L PBS. To detect caspase-2 activity, 90  $\mu$ g of the protein was combined with 200  $\mu$ M of Ac-VAVD-pNA in 150  $\mu$ L PBS. Reactions were incubated for 3 h at 37°C, and the activity was determined by measuring the absorbance at 405 nm.

### **Western blot analysis**

Cells were washed with PBS and lysed at 4°C with a lysis buffer containing 1% Triton X-100, protease inhibitor cocktail (Sigma-Aldrich), phosphatase inhibitor cocktail (Roche, Mannheim, Germany), and PBS. Lysates were clarified and the supernatants were subjected to western blotting as described previously (28).

### **Ethidium bromide (Et-Br) uptake assay and measurement**

Dye uptake experiments were performed as described previously (16). Briefly, THP-1 cells were seeded on glass coverslips and treated with 200 nM PMA for differentiation into

macrophages. Then, the cells were incubated with or without 1 mg/mL of TG for 24 h. Differentiated THP-1 cells were carefully washed three times with Dulbecco's phosphate buffered saline (DPBS) devoid of  $\text{Ca}^{2+}$  and loaded with pre-warmed DPBS containing Et-Br (final concentration 25  $\mu\text{g/mL}$ ) for 5 min at 37°C. The supernatant was discarded and cells washed twice with pre-warmed DPBS and fixed in 2% paraformaldehyde for 5 min. After fixation, cells were washed twice with DPBS and incubated with a DAPI stain solution (Vector Lab, Burlingame, CA, USA) in the dark for 10 min. Intracellular Et-Br fluorescence in cell monolayers was observed using a laser confocal scanning microscope (LSM 710, Zeiss, Heidenheim, Germany). Et-Br fluorescence intensity was measured using Image J (National Institute of Health, Bethesda, MD, USA).

### Statistical analysis

Statistical analysis was performed using GraphPad Prism 5 (GraphPad Software Inc., San Diego, CA, USA). The  $p$ -values were calculated using Student's  $t$ -test. Data are shown as the mean and standard error of the mean (SEM). Each experiment was conducted three times and the data were pooled for analysis. Differences were considered to be statistically significant at  $*p < 0.05$ ,  $**p < 0.01$ , or  $***p < 0.001$ .

### ACKNOWLEDGMENTS

This research was supported by the Korea Nazarene University Research Grant 2020.

### CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

**FIGURE LEGENDS****Figure 1. Potassium efflux is involved in TG-induced THP-1 macrophage cell death.**

(A) THP-1 cells were differentiated with PMA and incubated with or without TG for an additional 24 h. Extracellular potassium concentration was measured by atomic absorption spectrometry. (B) THP-1 macrophages were incubated with TG in the absence or presence of the potassium efflux inhibitor, glyburide. Viable cells were enumerated by the trypan blue dye exclusion assay. The number of viable cells in THP-1 macrophages without TG treatment was set as 100%. (C) THP-1 macrophages were incubated with TG in the presence of KCl for 24 h and the trypan blue exclusion assay was performed. (D) The cleaved form of PARP, and the cleavage of caspase-3, -7, -8 and -9, were detected by Western blotting. (E) The cleavage of caspase-1 was detected by Western blotting. (F) THP-1 macrophages were incubated with TG in the presence of glyburide for 24 h, after which caspase-1 activity was assessed. The absorbance of THP-1 macrophages without TG treatment was set as 100%. (G) THP-1 macrophages were incubated with TG in the presence of glyburide for 24 h and caspase-2 activity was assessed. All data are expressed as the mean  $\pm$  SEM of three independent experiments. P-values were determined with Student's t-test. \*  $P < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ .

**Figure-2. An increase in extracellular ATP induces TG-triggered macrophage cell death.**

(A) THP-1 macrophages were incubated with TG in the presence of ATP for 24 h and the trypan blue exclusion assay was performed. The number of viable cells in THP-1 macrophages without TG treatment was set as 100%. (B) The cleaved form of PARP was detected by Western blotting. (C) THP-1 macrophages were incubated with TG in the presence of ATP for 24 h, after which caspase-1 activity was assessed. The absorbance of THP-1 macrophages without treatment TG was set as 100%. (D) THP-1 macrophages were incubated with TG in



the presence of ATP for 24 h and caspase-2 activity was assessed. **(E)** THP-1 macrophages were incubated with TG in the presence of the ATP hydrolysis enzyme apyrase for 24 h, and viable cells were enumerated. The number of viable cells in THP-1 macrophages without TG treatment was set as 100%. **(F)** The cleaved form of PARP, and the cleavage of caspase-3, -7, -8, and -9 were detected by Western blotting. **(G)** THP-1 macrophages were incubated with TG in the presence of apyrase for 24 h, after which caspase-1 activity was assessed. The absorbance of THP-1 macrophages without TG treatment was set as 100%. **(H)** THP-1 macrophages were incubated with TG in the presence of apyrase for 24 h and caspase-2 activity was assessed. All data are expressed as the mean  $\pm$  SEM of three independent experiments. P-values were determined with Student's t-test. \*  $P < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ .

**Figure 3. TG-induced cell death involves activation of pannexin-1 channel in THP-1 macrophages.**

**(A)** THP-1 macrophages were incubated with the indicated concentration of TG for the indicated times. The cleaved form of pannexin-1 was detected by Western blotting. **(B)** THP-1 macrophages were incubated with or without TG for 24 h and an ethidium bromide uptake assay was performed to detect dye uptake by pannexin-1 channels (scale bars, 200  $\mu$ m). **(C)** Relative ethidium bromide intensity measured. The fluorescence intensity of THP-1 macrophages without TG treatment was set as 1.0. **(D)** THP-1 macrophages were incubated with TG in the presence of the pannexin-1 channel inhibitor probenecid for 24 h, then viable cells were enumerated. The number of viable cells in THP-1 macrophages without TG treatment was set as 100%. **(E)** The cleaved form of PARP, and the cleavage of caspase-3, -7, -8, and -9 were detected by Western blotting. **(F)** THP-1 macrophages were incubated with TG in the presence of probenecid for 24 h, after which caspase-1 activity was assessed. The absorbance of THP-1 macrophages without TG treatment was set as 100%. **(G)** THP-1

macrophages were incubated with TG in the presence of probenecid for 24 h and caspase-2 activity was assessed. All data are expressed as the mean  $\pm$  SEM of three independent experiments. P-values were determined with Student's t-test. \*  $P < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ .

**Figure 4. TG-triggered cell death is mediated by ATP released via activated pannexin-1 channels.**

(A) THP-1 macrophages were incubated with TG in the presence of probenecid and ATP for 24 h and viable cells were enumerated. The number of viable cells in THP-1 macrophages without TG treatment was set as 100%. (B) The cleaved form of PARP, and the cleavage of caspase-3, -7, -8, and -9 were detected by Western blotting. (C) THP-1 macrophages were incubated with TG in the presence of probenecid and ATP for 24 h, after which caspase-1 activity was assessed. The absorbance of THP-1 macrophages without TG treatment was set as 100%. (D) THP-1 macrophages were incubated with TG in the presence of probenecid and ATP for 24 h then caspase-2 activity was assessed. All data are expressed as the mean  $\pm$  SEM of three independent experiments. P-values were determined with Student's t-test. \*  $P < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ .

## REFERENCES

1. Lozano R, Naghavi M, Foreman K et al (2012) Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380, 2095-2128
2. Rafieian-Kopaei M, Setorki M, Douidi M, Baradaran A and Nasri H (2014) Atherosclerosis: process, indicators, risk factors and new hopes. *Int J Prev Med* 5, 927-946
3. Parthasarathy S, Raghavamenon A, Garelnabi MO and Santanam N (2010) Oxidized low-density lipoprotein. *Methods Mol Biol* 610, 403-417
4. Thomsen M, Varbo A, Tybjaerg-Hansen A and Nordestgaard BG (2014) Low nonfasting triglycerides and reduced all-cause mortality: a mendelian randomization study. *Clin Chem* 60, 737-746
5. Bersot T, Haffner S, Harris WS, Kellick KA and Morris CM (2006) Hypertriglyceridemia: management of atherogenic dyslipidemia. *J Fam Pract* 55, S1-8
6. Son SJ, Rhee K-J, Lim J, Kim TU, Kim T-J and Kim YS (2013) Triglyceride-induced macrophage cell death is triggered by caspase-1. *Biological and Pharmaceutical Bulletin* 36, 108-113
7. Lim J, Kim HK, Kim SH, Rhee KJ and Kim YS (2017) Caspase-2 mediates triglyceride (TG)-induced macrophage cell death. *BMB Rep* 50, 510-515
8. Park IS and Kim JE (2002) Potassium efflux during apoptosis. *J Biochem Mol Biol* 35, 41-46
9. Abdalah R, Wei L, Francis K and Yu SP (2006) Valinomycin-induced apoptosis in Chinese hamster ovary cells. *Neurosci Lett* 405, 68-73
10. Imre G, Heering J, Takeda AN et al (2012) Caspase-2 is an initiator caspase responsible for pore-forming toxin-mediated apoptosis. *The EMBO journal* 31, 2615-2628

11. Tinker A, Aziz Q and Thomas A (2014) The role of ATP-sensitive potassium channels in cellular function and protection in the cardiovascular system. *Br J Pharmacol* 171, 12-23
12. Coetzee WA (1992) ATP-sensitive potassium channels and myocardial ischemia: why do they open?. *Cardiovasc Drugs Ther* 6, 201-208
13. Ruan Z, Orozco IJ, Du J and Lu W (2020) Structures of human pannexin 1 reveal ion pathways and mechanism of gating. *Nature*. 584, 646-651
14. Bortner CD, Hughes FM, Jr. and Cidlowski JA (1997) A primary role for K<sup>+</sup> and Na<sup>+</sup> efflux in the activation of apoptosis. *J Biol Chem* 272, 32436-32442
15. Yu SP, Yeh CH, Gottron F, Wang X, Grabb MC and Choi DW (1999) Role of the outward delayed rectifier K<sup>+</sup> current in ceramide-induced caspase activation and apoptosis in cultured cortical neurons. *J Neurochem* 73, 933-941
16. Brough D, Pelegrin P and Rothwell NJ (2009) Pannexin-1-dependent caspase-1 activation and secretion of IL-1 $\beta$  is regulated by zinc. *Eur J Immunol* 39, 352-358
17. Bao L, Locovei S and Dahl G (2004) Pannexin membrane channels are mechanosensitive conduits for ATP. *FEBS Lett* 572, 65-68
18. Penuela S, Gehi R and Laird DW (2013) The biochemistry and function of pannexin channels. *Biochim Biophys Acta* 1828, 15-22
19. Tabas I (2009) Macrophage apoptosis in atherosclerosis: consequences on plaque progression and the role of endoplasmic reticulum stress. *Antioxid Redox Signal* 11, 2333-2339
20. Salvayre R, Auge N, Benoist H and Negre-Salvayre A (2002) Oxidized low-density lipoprotein-induced apoptosis. *Biochim Biophys Acta* 1585, 213-221
21. Giovannini C, Vari R, Scazzocchio B et al (2011) OxLDL induced p53-dependent apoptosis by activating p38MAPK and PKC $\delta$  signaling pathways in J774A.1

- macrophage cells. *J Mol Cell Biol* 3, 316-318
22. Klempfner R, Erez A, Sagit B-Z et al (2016) Elevated triglyceride level is independently associated with increased all-cause mortality in patients with established coronary heart disease: twenty-two-year follow-up of the bezafibrate infarction prevention study and registry. *Circulation: Cardiovascular Quality and Outcomes* 9, 100-108
  23. Chekeni FB, Elliott MR, Sandilos JK et al (2010) Pannexin 1 channels mediate 'find-me' signal release and membrane permeability during apoptosis. *Nature* 467, 863-867
  24. Schilling T and Eder C (2009) Lysophosphatidylcholine- and MCP-1-induced chemotaxis of monocytes requires potassium channel activity. *Pflügers Arch* 459, 71-77
  25. Ling MY, Ma ZY, Wang YY et al (2013) Up-regulated ATP-sensitive potassium channels play a role in increased inflammation and plaque vulnerability in macrophages. *Atherosclerosis* 226, 348-355
  26. Lim J, Kim SH, Kang YW et al (2014) Triglyceride up-regulates expression of ABCG1 in PMA-induced THP-1 macrophages through activation of JNK and p38 MAPK pathways. *Biomed Sci Letters* 20, 237-243
  27. Joo D, Woo JS, Cho KH et al (2016) Biphasic activation of extracellular signal-regulated kinase (ERK) 1/2 in epidermal growth factor (EGF)-stimulated SW480 colorectal cancer cells. *BMB Rep* 49, 220-225
  28. Jo HS, Kim DS, Ahn EH et al (2016) Protective effects of Tat-NQO1 against oxidative stress-induced HT-22 cell damage, and ischemic injury in animals. *BMB Rep* 49, 617-




Fig. 1.




Fig. 2.




Fig. 3.






Fig. 4.